

▼▲▼▲▼▲▼▲ 56:171 Operations Research ▼▲▼▲▼▲▼▲
 ▲▼▲▼▲▼▲▲ Final Examination Solutions ▲▼▲▼▲▼▲▲
 ▼▲▼▲▼▲▼▲ December 15, 1999 ▼▲▼▲▼▲▼▲

- Write your name on the first page, and initial the other pages.
- Answer both Parts A and B, and 4 (out of 5) problems from Part C.

		Possible
Part A:	Miscellaneous multiple choice	17
Part B:	Sensitivity analysis (LINDO)	14
Part C:	1. Discrete-time Markov chains I	15
	2. Discrete-time Markov chains II	15
	3. Continuous-time Markov chains	15
	4. Integer Programming Models	15
	5. Dynamic programming	<u>15</u>
	<i>total possible:</i>	<u>88</u>

▼▲▼▲▼▲▼ PART A ▼▲▼▲▼▲▼

Multiple Choice: Write the appropriate letter (a, b, c, d, or e) : (NOTA = None of the above).


- e 1. If, in the optimal *primal* solution of an LP problem (min cx st $Ax \geq b, x \geq 0$), there is zero slack in constraint #1, then in the optimal dual solution,
- a. dual variable #1 must be zero c. slack variable for dual constraint #1 must be zero
 b. dual variable #1 must be positive d. dual constraint #1 must be slack e. NOTA
- Note: Dual variable #1 may or may not be zero (if zero, solution is "degenerate").*
- c 2. If, in the optimal solution of the *dual* of an LP problem (min cx subject to: $Ax \geq b, x \geq 0$), dual variable #2 is positive, then in the optimal *primal* solution,
- a. variable #2 must be zero c. slack variable for constraint #2 must be zero
 b. variable #2 must be positive d. constraint #2 must be slack e. NOTA
- b 3. If $X_{ij} > 0$ in the transportation problem, then dual variables U and V *must* satisfy
- a. $C_{ij} > U_i + V_j$ c. $C_{ij} < U_i + V_j$ e. $C_{ij} = U_i - V_j$
 b. $C_{ij} = U_i + V_j$ d. $C_{ij} + U_i + V_j = 0$ f. NOTA
- c 4. For a discrete-time Markov chain, let P be the matrix of transition probabilities. The sum of each...
- a. column is 1 c. row is 1
 b. column is 0 d. row is 0 e. NOTA
- d 5. In a birth/death process model of a queue, the time between arrivals is assumed to
- a. have the Beta distribution c. be constant
 b. have the Normal distribution d. have the exponential distribution e. NOTA
- c 6. In an M/M/1 queue, if the arrival rate $= \lambda < \mu =$ service rate, then
- a. $\pi_0 = 1$ in steady state c. $\pi_i > 0$ for all i e. the queue is not a birth-death process
 b. no steady state exists d. $\pi_0 = 0$ in steady state f. NOTA
- d 7. If there is a tie in the "minimum-ratio test" of the simplex method, the solution in the next tableau
- a. will be nonbasic c. will have a worse objective value
 a. will be nonfeasible d. will be degenerate e. NOTA
- a 8. An absorbing state of a Markov chain is one in which the probability of
- a. moving out of that state is zero c. moving out of that state is one.
 b. moving into that state is one. d. moving into that state is zero e. NOTA

The problems (9)-(12) below refer to the following LP:

$$\begin{aligned}
 &\text{Maximize } 3X_1 + 5X_2 \\
 &\text{subject to } 2X_1 - X_2 \leq 5 \\
 &\quad X_1 + 3X_2 \leq 15 \\
 &\quad X_1 - 2X_2 \geq -4 \\
 &\quad X_1 \geq 0, X_2 \geq 0
 \end{aligned}$$

(with inequalities converted to equations:)

$$\begin{aligned}
 &\text{Maximize } 3X_1 + 5X_2 \\
 &\text{subject to } 2X_1 - X_2 + X_3 = 5 \\
 &\quad X_1 + 3X_2 + X_4 = 15 \\
 &\quad X_1 - 2X_2 - X_5 = -4 \\
 &\quad X_j \geq 0, j=1,2, 3,4,5
 \end{aligned}$$



- a or b 9. The feasible region includes (possibly with others) points:
- | | | |
|--------------|--------------|----------------|
| a. A, B, & C | c. D, C, & E | e. D, E, & G |
| b. B & F | d. E, F, & G | f. <i>NOTA</i> |
- e 10. At point **F**, the basic variables include the variables
- | | | |
|-----------------------|-----------------------|-----------------------|
| a. X_1, X_2 & X_3 | c. X_2, X_4 & X_5 | e. X_1, X_2 & X_5 |
| b. X_1, X_3 & X_4 | d. X_1, X_2 & X_4 | f. <i>NOTA</i> |
- f 11. Which point is degenerate in the primal problem?
- | | | |
|------------|------------|----------------|
| a. point A | c. point C | e. point E |
| b. point B | d. point D | f. <i>NOTA</i> |
- a 12. The *dual* of this LP has the following constraints (not including nonnegativity or nonpositivity):
- | | |
|-------------------------------------|--|
| a. 2 constraints of type (\geq) | d. one each of type \geq & $=$ |
| b. one each of type \leq & \geq | e. 2 of type \leq and 1 of type \geq |
| c. 2 constraints of type (\leq) | f. <i>NOTA</i> |
- d 13. The dual of the LP has the following types of variables:
- | | |
|---|---------------------------------|
| a. three non-negative variables | e. three non-positive variables |
| b. one non-negative and two non-positive variables | f. <i>NOTA</i> |
| c. two non-negative variables and one unrestricted in sign | |
| d. two non-negative variables and one non-positive variable | |
- d 14. If point **F** is optimal, then which dual variables must be zero, according to the *Complementary Slackness Theorem* ?
- | | |
|--------------------|---------------|
| a. Y_1 and Y_2 | d. Y_1 only |
| b. Y_1 and Y_3 | e. Y_2 only |
| c. Y_2 and Y_3 | f. Y_3 only |
15. The number of basic variables in a solution of a transportation problem with 5 sources and 7 destinations is 11
- d 16. A balanced transportation problem is one in which
- | | | |
|--------------------------------|------------------------------------|----------------|
| a. # sources = # destinations | c. supplies & demands all 1 | e. <i>NOTA</i> |
| b. cost coefficients are all 1 | d. sum of supplies = sum of demand | |
- c 17. An assignment problem is a transportation problem for which
- | | | |
|--------------------------------|------------------------------------|----------------|
| a. # sources = # destinations | c. supplies & demands all 1 | e. <i>NOTA</i> |
| b. cost coefficients are all 1 | d. sum of supplies = sum of demand | |

▼▲▼▲▼▲▼ PART B ▼▲▼▲▼▲▼

Sensitivity Analysis in LP.

Zales Jewelers uses rubies and sapphires to produce two types of rings. A type 1 ring requires 2 rubies, 3 sapphires, and 1 hour of jeweler's labor. A type 2 ring requires 3 rubies, 2 sapphires, and 2 hours of jeweler's labor. Each type 1 ring sells for \$400, and each type 2 ring sells for \$500. All rings produced by Zales can be sold. At present, Zales has 100 rubies, 120 sapphires, and 70 hours of jeweler's labor available. Extra rubies can be purchased at a cost of \$100 each. Market demand requires that the company produce at least 20 type 1 rings and at least 25 type 2 rings. To maximize profit, Zales should solve the following LP:

X1= type 1 rings produced.

X2= type 2 rings produced

R = number of rubies purchased.

$$\text{MAX } z=400X1 + 500X2 - 100R$$

$$\text{s.t. } 2X1 + 3X2 - R \leq 100$$

$$3X1 + 2X2 \leq 120$$

$$X1 + 2X2 \leq 70$$

$$X1 \geq 20$$

$$X2 \geq 25$$

$$X10, X20, R0$$

The LINDO output for this problem follows:

```

MAX      400 X1 + 500 X2 - 100 R
SUBJECT TO
    2)    2 X1 + 3 X2 - R <=   100
    3)    3 X1 + 2 X2 <=   120
    4)    X1 + 2 X2 <=    70
    5)    X1 >=    20
    6)    X2 >=    25

END

      OBJECTIVE FUNCTION VALUE
1)    19000.00

      VARIABLE            VALUE            REDUCED COST
      X1                   20.000000            .000000
      X2                   25.000000            .000000
      R                    15.000000            .000000

      ROW    SLACK OR SURPLUS    DUAL PRICES
      2)           .000000           100.000000
      3)          10.000000            .000000
      4)           .000000           200.000000
      5)           .000000            .000000
      6)          -.000000          -200.000000

```

RANGES IN WHICH THE BASIS IS UNCHANGED:

VARIABLE	CURRENT COEF	OBJ COEFFICIENT RANGES	
		ALLOWABLE INCREASE	ALLOWABLE DECREASE
X1	400.000000	INFINITY	100.000000
X2	500.000000	200.000000	INFINITY
R	-100.000000	100.000000	100.000000

ROW	CURRENT RHS	RIGHTHAND SIDE RANGES	
		ALLOWABLE INCREASE	ALLOWABLE DECREASE
2	100.000000	15.000000	INFINITY
3	120.000000	INFINITY	10.000000
4	70.000000	3.333333	.000000
5	20.000000	.000000	INFINITY
6	25.000000	.000000	2.500000

Solutions

semester, a 15% chance of remaining a junior, and a 5% chance of quitting. (We will assume that a student who quits never re-enrolls.)

Powers of P:

$$P^2 = \begin{array}{c|cccccc} & \backslash & 1 & 2 & 3 & 4 & 5 & 6 \\ \hline m & & & & & & & \\ \hline 1 & 0.01 & 0.16 & 0.68 & 0 & 0.15 & 0 & \\ \hline 2 & 0 & 0.01 & 0.187 & 0.68 & 0.123 & 0 & \\ \hline 3 & 0 & 0 & 0.0144 & 0.176 & 0.1296 & 0.68 & \\ \hline 4 & 0 & 0 & 0 & 0.01 & 0.055 & 0.935 & \\ \hline 5 & 0 & 0 & 0 & 0 & 1 & 0 & \\ \hline 6 & 0 & 0 & 0 & 0 & 0 & 1 & \end{array}$$

$$P^3 = \begin{array}{c|cccccc} & \backslash & 1 & 2 & 3 & 4 & 5 & 6 \\ \hline m & & & & & & & \\ \hline 1 & 0.001 & 0.024 & 0.2176 & 0.544 & 0.2134 & 0 & \\ \hline 2 & 0 & 0.001 & 0.03094 & 0.2176 & 0.17246 & 0.578 & \\ \hline 3 & 0 & 0 & 0.001728 & 0.02912 & 0.139552 & 0.8296 & \\ \hline 4 & 0 & 0 & 0 & 0.001 & 0.0555 & 0.9435 & \\ \hline 5 & 0 & 0 & 0 & 0 & 1 & 0 & \\ \hline 6 & 0 & 0 & 0 & 0 & 0 & 1 & \end{array}$$

$$P^4 = \begin{array}{c|cccccc} & \backslash & 1 & 2 & 3 & 4 & 5 & 6 \\ \hline from & & & & & & & \\ \hline 1 & 0.0001 & 0.0032 & 0.046512 & 0.22848 & 0.259308 & 0.4624 & \\ \hline 2 & 0 & 0.0001 & 0.0045628 & 0.046512 & 0.185865 & 0.76296 & \\ \hline 3 & 0 & 0 & 0.00020736 & 0.0042944 & 0.141146 & 0.854352 & \\ \hline 4 & 0 & 0 & 0 & 0.0001 & 0.05555 & 0.94435 & \\ \hline 5 & 0 & 0 & 0 & 0 & 1 & 0 & \\ \hline 6 & 0 & 0 & 0 & 0 & 0 & 1 & \end{array}$$

Absorption Probabilities A:

	\	5	6
from		-----	-----
1		0.279212	0.720788
2		0.189113	0.810887
3		0.141414	0.858586
4		0.0555556	0.944444

Expected # of Visits E=

	\	1	2	3	4
from		-----	-----	-----	-----
1		1.11111	0.987654	0.953984	0.847986
2		0	1.11111	1.07323	0.953984
3		0	0	1.13636	1.0101
4		0	0	0	1.11111

Select the **nearest** available numerical choice in answering the questions below.

- e 1. The number of *transient* states in this Markov chain model is
 a. 0 c. 2 e. 4 g. *NOTA*
 b. 1 d. 3 f. 5
- c 2. The number of *absorbing* states in this Markov chain model is
 a. 0 c. 2 e. 4 g. *NOTA*
 b. 1 d. 3 f. 5
- c 3. The number of *recurrent* states in this Markov chain model is
 a. 0 c. 2 e. 4 g. *NOTA*
 b. 1 d. 3 f. 5
4. The closed sets of states in this Markov chain model are (circle all that apply!)
 a. {1} d. {4} g. {1,2,3,4} j. {2,3,4}
 b. {2} **e. {5}** h. {1,2,3,4} k. {3,4}
 c. {3} **f. {6}** **i. {5,6}** l. **{1,2,3,4,5,6}**

5. The *minimal* closed sets of states in this Markov chain model are (circle all that apply!)

- | | | | |
|---------|----------------|---------------|-------------------|
| a. {1 } | d. {4 } | g. {1,2,3,4 } | j. {2,3,4 } |
| b. {2 } | <u>e. {5 }</u> | h. {1,2,3,4 } | k. {3,4 } |
| c. {3 } | <u>f. {6 }</u> | i. {5,6 } | l. {1,2,3,4,5,6 } |

Suppose that at the beginning of the Fall '99 semester, Joe Cool was a *Freshman*.

g 6. What is the probability that Joe is a junior in Fall 2001? (choose nearest answer)

- | | | | | |
|--------|--------|--------|--------|---------|
| a. 10% | c. 30% | e. 50% | g. 70% | i. 90% |
| b. 20% | d. 40% | f. 60% | h. 80% | j. 100% |

e 7. What is the probability that Joe is a senior in Fall 2002? (choose nearest answer)

- | | | | | |
|--------|--------|--------|--------|---------|
| a. 10% | c. 30% | e. 50% | g. 70% | i. 90% |
| b. 20% | d. 40% | f. 60% | h. 80% | j. 100% |

* 8. What is the probability that Joe *first* becomes a senior in 2003? (choose nearest answer)

- | | | | | |
|--------|--------|--------|--------|--------|
| a. 5% | c. 15% | e. 25% | g. 35% | i. 45% |
| b. 10% | d. 20% | f. 30% | h. 40% | j. 50% |

* *This probability cannot be readily determined from the information given-- requires recursive computation of the first-passage probabilities.*

g 9. What is the probability that Joe *eventually* graduates? (choose nearest answer)

- | | | | | |
|--------|--------|--------|--------|---------|
| a. 10% | c. 30% | e. 50% | g. 70% | i. 90% |
| b. 20% | d. 40% | f. 60% | h. 80% | j. 100% |

c 10. What is the expected length of his academic career, in years? (choose nearest answer)

- | | | | | |
|---------------|---------------|---------------|---------------|----------------------|
| a. 3 year | c. 4 years | e. 4.5 years | g. 5 years | i. 5.5 years |
| b. 3.75 years | d. 4.25 years | f. 4.75 years | h. 5.25 years | j. ≥ 5.75 years |

e 11. What fraction of students graduate in *exactly* four years? (choose nearest answer)

- | | | | | | | |
|----------------|--------|--------|--------|--------|--------|----------------|
| a. $\leq 25\%$ | c. 35% | e. 45% | g. 55% | i. 65% | k. 75% | m. 85% |
| b. 30% | d. 40% | f. 50% | h. 60% | j. 70% | l. 80% | n. $\geq 90\%$ |

2. **Discrete-time Markov Chains II:** On New Year's Eve (December 31) of each year I determine whether my car is in good, fair, or broken-down condition. If my car is broken-down, I replace it on January 1st with a good used car.


- A good car will be good at the end of next year with probability 80%, fair with probability 15%, or broken-down with probability 5%.
- A fair car will be fair at the end of the next year with probability 50%, or broken-down with probability 50%.
- It costs \$10,000 to purchase a good used car; a fair car can be traded in for \$3000; and a broken-down car can be sold as junk for \$500.
- It costs \$1000 per year to operate a good car and \$1500 to operate a fair car. If a car breaks down during a year, the operating cost averages \$2000.

Assume that the cost of operating a car during a year depends on the type of car on hand at the *beginning* of the year, and that any break-down occurs only at the *end* of a year.

My policy is to drive my car until it breaks down, at which time I replace it with a good used car. Define a Markov chain model representing the condition of the car which I own on Dec. 31, with three states:

1. Good condition
2. Fair condition
3. Broken-down

The diagram below indicates the transition probabilities:



Below are the 2nd, 3rd, & 4th powers of P:

P ² =			P ³ =			P ⁴ =					
from \	1	2	3	from \	1	2	3	from \	1	2	3
1	0.6375	0.2025	0.16	1	0.598125	0.220875	0.181	1	0.584344	0.227306	0.18835
2	0.375	0.325	0.3	2	0.50625	0.26375	0.23	2	0.552188	0.242313	0.2055
3	0.6375	0.2025	0.16	3	0.598125	0.220875	0.181	3	0.584344	0.227306	0.18835

Mean 1st passage times

from \	1	2	3
1	1.73333	6.66667	5.2
2	3.73333	4.33333	2
3	1.73333	6.66667	5.2

_____ Which **one or more** equations must be satisfied by the steady state probabilities π_1 , π_2 , & π_3 ?

- a. $\pi_1 + \pi_2 + \pi_3 = 1$
- b. $\pi_1 + \pi_2 + \pi_3 = 0$
- c. $0.8\pi_1 + 0.8\pi_3 = \pi_1$
- d. $0.8\pi_1 + 0.15\pi_2 + 0.05\pi_3 = \pi_1$
- e. $0.8\pi_1 + 0.15\pi_2 + 0.05\pi_3 = \pi_3$
- f. $0.05\pi_1 + 0.5\pi_2 + 0.05\pi_3 = \pi_3$
- g. $0.8\pi_1 + 0.8\pi_3 = 0$
- h. $0.8\pi_1 + 0.15\pi_2 + 0.05\pi_3 = 0$

Write the expression which represents my average cost per year:

 \$1000 π_1 + \$1500 π_2 + (\$2000 + 10000 - 500) π_3

I should expect to replace my car once every 5.2 years.

If my current car is in *fair* condition, I should expect to replace it in 2 years.

 a 7. The number of transient states in this Markov chain is

- a. 0
- b. 1
- c. 2
- d. 3
- e. 4
- f. 5
- g. *NOTA*


 d 8. The number of recurrent states in this Markov chain is

- a. 0
- b. 1
- c. 2
- d. 3
- e. 4
- f. 5
- g. *NOTA*

3. Birth/Death Model of a Queue:


For each birth/death process below, pick the classification of the queue and write it in the blank to the left:

- a. M/M/1
- b. M/M/2
- c. M/M/4
- d. M/M/4/4
- e. M/M/2/2
- f. M/M/2/4/4
- g. M/M/2
- h. M/M/1/4
- i. M/M/1/4/4
- j. M/M/2/4
- k. M/M/4/2
- l. *NOTA*




 M/M/2/4/4 1.


Solutions




M/M/2/4 2.



M/M/2 3.




M/M/4 4.



M/M/1 5.

A job shop has four numerically controlled machines that are capable of operating on their own (i.e., without a human operator) once they have been set up with the proper cutting tools and all adjustments are made. Each setup requires the skills of an experienced machine operator, and the time need to complete a setup is exponentially distributed with a mean of 30 minutes. When the setup is complete, the machine operator pushes a button, and the machine requires no further attention until it has finished its job, when it is ready for another setup. The job times are exponentially distributed with a mean of one hour. Two operators have been assigned to this group of machines. (If only one machine requires attention, only one operator will tend it, rather than both working together.)

6. Indicate the transition rates on the diagram:



7. Write the expression which is used to evaluate π_0 :

$$\frac{1}{\pi_0} = 1 + \frac{4}{2} + \frac{4}{2} \times \frac{3}{4} + \frac{4}{2} \times \frac{3}{4} \times \frac{2}{4} + \frac{4}{2} \times \frac{3}{4} \times \frac{2}{4} \times \frac{1}{4}$$

The steady-state probability distribution is:

i	π_i	CDF
0	0.1839	0.1839
1	0.3678	0.5517
2	0.2759	0.8276
3	0.1379	0.9655
4	0.0345	1

c 8. What is the percent of the time that both machinists are idle? (Choose nearest value.)

- a. 10%
- b. 15%
- c. 20%
- d. 25%
- e. 30%
- f. 35%
- g. 40%
- h. 45% or more

e 9. What is the average number of machines in operation? (Choose nearest value.)

- a. 0.5
- b. 1.0
- c. 1.5
- d. 2.0
- e. 2.5 (2.52)
- f. 3.0
- g. 3.5
- h. 4.0

d 10. What is the utilization of each machine, i.e., the percent of the time that each machine is busy? (Choose nearest value.)

- a. $\leq 50\%$
- b. 55%
- c. 60%
- d. 65% (63.2%)
- e. 70%
- f. 75%
- g. 80%
- h. 85%
- i. 90%
- j. 95% or more

Suppose that the average rate at which machines complete jobs is 2.5/hour

c_11. What is the average length of the time interval between a machine's completion of a job and the starting of another job (in hours)? (Choose nearest value.)

- a. 0.5 c. 0.6 e. 0.7 g. 0.8 i. 0.9
 b. 0.55 d. 0.65 f. 0.75 h. 0.85 j. 0.95 or more

Little's Law implies that $L = \lambda W$ where $L = 0\pi_0 + 1\pi_1 + 2\pi_2 + 3\pi_3 + 4\pi_4$ is the average number in the queueing system. (Or L could be calculated as 4 minus your answer in (9) above. Hence $L = 4 - 2.52 = 1.48$ and $W = \frac{L}{\lambda} = \frac{1.48}{2.52/\text{hr}} = 0.592$ hr.

4. Integer Programming Models

The board of directors of a large manufacturing firm is considering the set of investments shown below: Let R_i be the annual revenue (in \$millions) from investment i and C_i the cost (in \$millions) to make investment i . The board wishes to maximize total annual revenues and invest no more than a total of 50 million dollars.

Investment i	Revenue R_i	Cost C_i	Condition
1	1	5	None
2	2	8	Only if #1
3	3	12	None
4	4	18	Must if #1 and #2
5	5	24	Not if both #3 and #4
6	6	27	None
7	7	30	Only if both #3 and #6

Define variables:

$X_i = 1$ if investment i is selected, else 0.

1. Formulate this problem without the "side conditions" as an integer LP.

$$\begin{aligned} & \text{Max } \sum_{i=1}^7 C_i X_i \\ & \text{s.t. } \sum_{i=1}^7 R_i X_i \leq 50 \\ & X_i \in \{0,1\} \text{ for } i = 1,2,\dots,7 \end{aligned}$$

This is an example of a special class of integer programming problems called *knapsack* problems.

2. Add a constraint or constraints to enforce the condition "Investment #2 can be selected only if #1 is selected".

$$X_2 \leq X_1$$

3. Add a constraint or constraints to enforce the condition "Investment #4 must be selected if both #1 & #2 are selected".

$$X_4 \geq X_1 + X_2 - 1$$

4. Add a constraint or constraints to enforce the condition "Investment #5 cannot be selected if both #3 & #4 are selected".

$$X_1 + X_2 + X_5 \leq 2$$

5. Add a constraint or constraints to enforce the condition "Investment #7 only if both #3 and #6 are selected".


Can use the pair of constraints

$$\begin{cases} X_7 \leq X_3 \\ X_7 \leq X_6 \end{cases}$$

or (summing the two constraints) $2X_7 \leq X_3 + X_6$

5. Optimization of System Reliability by DP: A system consists of 4 devices, each subject to possible failure, such that the system fails if any one or more of the devices fail:

Device	Reliability	Weight (kg)
1	75%	2
2	90%	3
3	80%	1
4	85%	2



Suppose that redundant units of devices 1 and 3 are included as shown above. (That is, system failure occurs if all 3 of device 1, or both of device 3, or device 2, or device 4 were to fail.)

d_1. The reliability of **device 3** in the system is:

- a. $.2^2 = 4\%$
- b. $.8^2 = 64\%$
- c. $1 - e^{-2 \times 0.2} = 32.97\%$
- d. $1 - .2^2 = 96\%$
- e. $1 - e^{-2 \times 0.8} = 79.8\%$
- f. $1 - .8^2 = 36\%$
- g. *NOTA*


c_2. The reliability of this entire **system** is:

- a. $1 - (0.75^3)(0.9)(0.8^2)(0.85) = 79.35\%$
- b. $(1 - e^{-3 \times 0.75})(1 - e^{-0.9})(1 - e^{-2 \times 0.8})(1 - e^{-0.85}) = 24.26\%$
- c. $(1 - 0.25^3)(1 - 0.1)(1 - 0.2^2)(1 - 0.15) = 72.29\%$
- d. $(1 - 0.75^3)(1 - 0.9)(1 - 0.8^2)(1 - 0.85) = 0.3\%$
- e. $(1 - e^{-3 \times 0.25})(1 - e^{-0.1})(1 - e^{-2 \times 0.2})(1 - e^{-0.15}) = 0.23\%$
- f. $1 - (1 - 0.25^3)(1 - 0.1)(1 - 0.2^2)(1 - 0.15) = 27.71\%$
- g. *NOTA*

b_3. The weight of this system is:

- a. 10 kg.
- b. 13 kg.
- c. 11 kg.
- d. 14 kg.
- e. 12 kg.
- f. none of the above

Suppose that we wish to find the system design having maximum reliability subject to a limit of 14 kg. weight.



c_4. Define a dynamic programming model with optimal value function $f_n(S)$, where $f_n(S)$ is

- a. the reliability of S redundant units of device #n.
- b. the maximum reliability of the system if n redundant units are allowed.
- c. the maximum reliability of devices 1 through n, if S kg. of weight is allocated to them.
- d. the maximum reliability of devices n through 4, if S kg. of weight is allocated to them.

c, f_5. The value of S_4 is (choose one or more!):

- a. the safety factor for device 4
- b. 15%
- c. the state of the DP system at stage 4
- d. 1 kg.
- e. the reliability of device #4
- f. 14 kg.

The following table shows the reliability of a device for various numbers of redundant units:

	# units		
Device	1	2	3
1	.75	.9375	.984375
2	.9	.99	.999
3	.8	.96	.992
4	.85	.9775	.996625

The following output is obtained during the solution of the DP model, where several values have been omitted. Note that the value -99.99999 is used to indicate -∞, i.e., an infeasible combination of s & x .

6. Enter the missing value for each of the entries in the tables:

α : $1 - 0.252 = \underline{0.9375}$ (see table above).

β : 0.92813 (maximum value in row of table)

γ : 2 (value of X in which β appears)

δ : $10 - 2 \times 3 = \underline{4}$ (remaining capacity after 2 units of weight 3 have been added to system)

ϵ : $0.992 \times f_2(12 - 3 \times 1) = 0.992 \times f_2(9) = 0.992 \times 0.88594 = \underline{0.87885}$

7. The optimal design, weighing 14 kg., has reliability: $f_4(14) = \underline{0.81817}$

8. The optimal design has 2 units of #1, 1 unit of #2, 3 units of #3, and 2 units of #4.

---Stage 1---

s \ x:	1	2	3	$f_1(s)$	optimal x	next state
2	0.75000	-99.99999	-99.99999	0.75000	1	0
3	0.75000	-99.99999	-99.99999	0.75000	1	1
4	0.75000	<u>α</u>	-99.99999	0.93750	2	0
5	0.75000	0.93750	-99.99999	0.93750	2	1
7	0.75000	0.93750	0.98438	0.98438	3	1
8	0.75000	0.93750	0.98438	0.98438	3	2
9	0.75000	0.93750	0.98438	0.98438	3	3
10	0.75000	0.93750	0.98438	0.98438	3	4
11	0.75000	0.93750	0.98438	0.98438	3	5
12	0.75000	0.93750	0.98438	0.98438	3	6
13	0.75000	0.93750	0.98438	0.98438	3	7
14	0.75000	0.93750	0.98438	0.98438	3	8

---Stage 2---

s \ x:	1	2	3	$f_1(s)$	optimal x	next state
5	0.67500	-99.99999	-99.99999	0.67500	1	2
6	0.67500	-99.99999	-99.99999	0.67500	1	3
7	0.84375	-99.99999	-99.99999	0.84375	1	4
8	0.84375	0.74250	-99.99999	0.84375	1	5
9	0.8859	0.74250	-99.99999	0.88594	1	6
10	0.88594	0.92813	-99.99999	<u>β</u>	<u>γ</u>	<u>δ</u>
11	0.88594	0.92813	0.74925	0.92813	2	5
12	0.88594	0.97453	0.74925	0.97453	2	6
13	0.88594	0.97453	0.93656	0.97453	2	7
14	0.88594	0.97453	0.93656	0.97453	2	8

Solutions

---Stage 3---

s \ x:				f ₁ (s)	optimal	next
	1	2	3		x	state
6	0.54000	-99.99999	-99.99999	0.54000	1	5
7	0.54000	0.64800	-99.99999	0.64800	2	5
8	0.67500	0.64800	0.66960	0.67500	1	7
9	0.67500	0.81000	0.66960	0.81000	2	7
10	0.70875	0.81000	0.83700	0.83700	3	7
11	0.74250	0.85050	0.83700	0.85050	2	9
12	0.74250	0.89100	$\underline{\underline{\epsilon}}$	0.89100	2	10
13	0.77963	0.89100	0.92070	0.92070	3	10
14	0.77963	0.93555	0.92070	0.93555	2	12

---Stage 4---

s \ x:				f ₁ (s)	optimal	next
	1	2	3		x	state
8	0.45900	-99.99999	-99.99999	0.45900	1	6
9	0.55080	-99.99999	-99.99999	0.55080	1	7
10	0.57375	0.52785	-99.99999	0.57375	1	8
11	0.68850	0.63342	-99.99999	0.68850	1	9
12	0.71145	0.65981	0.53818	0.71145	1	10
13	0.72293	0.79178	0.64581	0.79178	2	9
14	0.75735	0.81817	0.67272	0.81817	2	10