

# DUALITY IN LINEAR PROGRAMMING

This hypercard stack prepared by:  
Dennis Bricker  
Dept of I.E., U. of Iowa  
dennis-bricker@uiowa.edu

## Outline of Contents

- Definition of Symmetric Primal/Dual Pair
- Writing the dual of a general Primal LP
- Weak Duality Theorem
- Simplex multipliers solve the dual!
- Economic interpretation of dual LP
- Fundamental Duality Theorem
- Farkas' Lemma
- Complementary Slackness

The **SYMMETRIC** primal/dual pair:

Primal:

Minimize  $c^t x$   
subject to:

$$\begin{aligned} Ax &\geq b \\ x &\geq 0 \end{aligned}$$

Dual:

Maximize  $b^t y$   
subject to:

$$\begin{aligned} A^t y &\leq c \\ y &\geq 0 \end{aligned}$$

where  $A$  is an  $m \times n$  matrix,  $x$  &  $c$  are vectors of length  $n$ ,  
and  $y$  &  $b$  are vectors of length  $m$ . (Note:  $A^t$  denotes transpose  
of the matrix  $A$ .)

Note the following characteristics:

- the primal LP is  $m \times n$ , i.e.,  $m$  constraints (not including nonnegativity) and  $n$  variables
- the dual LP is  $n \times m$ , i.e.,  $n$  constraints (not including nonnegativity) and  $m$  variables

Primal:

minimize  $c^t x$   
subject to:

$$\begin{aligned} Ax &\geq b \\ x &\geq 0 \end{aligned}$$

Dual:

Maximize  $b^t y$   
subject to:

$$\begin{aligned} A^t y &\leq c \\ y &\geq 0 \end{aligned}$$

Note the following characteristics:

- for every variable in the primal problem, there is a corresponding inequality constraint in the dual problem
- for every inequality constraint (not including nonnegativity), there is a corresponding dual variable

Primal:

$$\begin{aligned} &\text{minimize } c^t x \\ &\text{subject to:} \\ &\quad Ax \geq b \\ &\quad x \geq 0 \end{aligned}$$

Dual:

$$\begin{aligned} &\text{Maximize } b^t y \\ &\text{subject to:} \\ &\quad A^t y \leq c \\ &\quad y \geq 0 \end{aligned}$$

Note the following characteristics:


- the right-hand-side vector ( $b$ ) of the primal problem serves as the objective function coefficient vector of the dual problem.

Primal:

$$\begin{aligned} &\text{minimize } c^t x \\ &\text{subject to:} \\ &\quad Ax \geq b \\ &\quad x \geq 0 \end{aligned}$$

Dual:

$$\begin{aligned} &\text{Maximize } b^t y \\ &\text{subject to:} \\ &\quad A^t y \leq c \\ &\quad y \geq 0 \end{aligned}$$



## Example

*Primal*

$$\text{Minimize } 20x_1 + 10x_2$$

subject to:

$$5x_1 + x_2 \geq 6$$

$$2x_1 + 2x_2 \geq 8$$

$$x_1 \geq 0, x_2 \geq 0$$

*Dual*

$$\text{Maximize } 6y_1 + 8y_2$$

subject to:

$$5y_1 + 2y_2 \leq 20$$

$$y_1 + 2y_2 \leq 10$$

$$y_1 \geq 0, y_2 \geq 0$$

*Primal*

$$\underline{\text{Minimize } 20x_1 + 10x_2}$$

subject to:

$$5x_1 + x_2 \geq 6$$

$$2x_1 + 2x_2 \geq 8$$

$$x_1 \geq 0, x_2 \geq 0$$

*Dual*

$$\underline{\text{Maximize } 6y_1 + 8y_2}$$

subject to:

$$5y_1 + 2y_2 \leq 20$$

$$y_1 + 2y_2 \leq 10$$

$$y_1 \geq 0, y_2 \geq 0$$

**The primal problem is a MINIMIZATION with  $\geq$  constraints, while the dual problem is a MAXIMIZATION with  $\leq$  constraints!**

*Primal**Dual*

$$\text{Minimize } 20x_1 + 10x_2$$

$$\text{Maximize } 6y_1 + 8y_2$$

subject to:

subject to:

$$5x_1 + x_2 \geq 6$$

$$5y_1 + 2y_2 \leq 20$$

$$2x_1 + 2x_2 \geq 8$$

$$y_1 + 2y_2 \leq 10$$

$$x_1 \geq 0, x_2 \geq 0$$

$$y_1 \geq 0, y_2 \geq 0$$

**The objective coefficients of the primal serve as the right-hand-side of the dual problem!**

*Primal**Dual*

$$\text{Minimize } 20x_1 + 10x_2$$

$$\text{Maximize } 6y_1 + 8y_2$$

subject to:

subject to:

$$5x_1 + x_2 \geq 6$$

$$5y_1 + 2y_2 \leq 20$$

$$2x_1 + 2x_2 \geq 8$$

$$y_1 + 2y_2 \leq 10$$

$$x_1 \geq 0, x_2 \geq 0$$

$$y_1 \geq 0, y_2 \geq 0$$

**... and conversely, the right-hand-side of the primal problem serves as objective coefficients of the dual problem!**

*Primal*Minimize  $20x_1 + 10x_2$ 

subject to:

$$5x_1 + x_2 \geq 6$$

$$2x_1 + 2x_2 \geq 8$$

$$x_1 \geq 0, x_2 \geq 0$$

*Dual*Maximize  $6y_1 + 8y_2$ 

subject to:

$$5y_1 + 2y_2 \leq 20$$

$$y_1 + 2y_2 \leq 10$$

$$y_1 \geq 0, y_2 \geq 0$$

**To every constraint in the primal, there corresponds a dual variable.**

*Primal*Minimize  $20x_1 + 10x_2$ 

subject to:

$$5x_1 + x_2 \geq 6$$

$$2x_1 + 2x_2 \geq 8$$

$$x_1 \geq 0, x_2 \geq 0$$

*Dual*Maximize  $6y_1 + 8y_2$ 

subject to:

$$5y_1 + 2y_2 \leq 20$$

$$y_1 + 2y_2 \leq 10$$

$$y_1 \geq 0, y_2 \geq 0$$

**To every variable in the primal problem, there corresponds a constraint in the dual problem.**

*Primal*Minimize  $20x_1 + 10x_2$ 

subject to:

$$5x_1 + x_2 \geq 6$$

$$2x_1 + 2x_2 \geq 8$$

$$x_1 \geq 0, x_2 \geq 0$$

*Dual*Maximize  $6y_1 + 8y_2$ 

subject to:

$$5y_1 + 2y_2 \leq 20$$

$$y_1 + 2y_2 \leq 10$$

$$y_1 \geq 0, y_2 \geq 0$$

**Both primal and dual problems include nonnegativity constraints on the variables.**

*Suppose that we have an inequality reversed in the primal problem, for example:*

Minimize  $20x_1 + 10x_2$ 

subject to:

$$5x_1 + x_2 \leq 6$$

$$2x_1 + 2x_2 \geq 8$$

$$x_1 \geq 0, x_2 \geq 0$$

*note the reversed direction!*

*How do we write the dual of this problem?*

*First we must transform the problem:*

**We multiply the offending inequality by  $-1$ , thereby reversing the direction of the inequality:**

$$\text{Minimize } 20x_1 + 10x_2$$

$$\text{Minimize } 20x_1 + 10x_2$$

subject to:

subject to:

$$5x_1 + x_2 \leq 6 \implies -5x_1 - x_2 \geq -6$$

$$2x_1 + 2x_2 \geq 8$$

$$2x_1 + 2x_2 \geq 8$$

$$x_1 \geq 0, x_2 \geq 0$$

$$x_1 \geq 0, x_2 \geq 0$$

**Now the problem is in the form of the primal in the symmetric primal/dual pair. We can therefore write its DUAL problem:**

$$\text{Minimize } 20x_1 + 10x_2$$

$$\text{Maximize } -6y_1 + 8y_2$$

subject to:

subject to

$$-5x_1 - x_2 \geq -6$$

$$-5y_1 + 2y_2 \leq 20$$

$$2x_1 + 2x_2 \geq 8$$

$$-y_1 + 2y_2 \leq 10$$

$$x_1 \geq 0, x_2 \geq 0$$

$$y_1 \geq 0, y_2 \geq 0$$

*It is interesting to now make a change of variable: let  $y'_1 = -y_1$*


Maximize  $-6y_1 + 8y_2$

subject to

$$-5y_1 + 2y_2 \leq 20$$

$$-y_1 + 2y_2 \leq 10$$

$$y_1 \geq 0, y_2 \geq 0$$



Maximize  $6y'_1 + 8y_2$

subject to

$$5y'_1 + 2y_2 \leq 20$$

$$y'_1 + 2y_2 \leq 10$$

$$y'_1 \leq 0, y_2 \geq 0$$



*Same as dual of the symmetric primal/dual pair, except for non-positivity replacing non-negativity!*

*Suppose that, rather than an inequality constraint, we had an equality constraint. For example:*

Minimize  $20x_1 + 10x_2$

subject to:

$$5x_1 + x_2 = 6$$

$$2x_1 + 2x_2 \geq 8$$

$$x_1 \geq 0, x_2 \geq 0$$

*What is its DUAL problem?*

*We must first transform the equality constraint into equivalent inequalities:*

$$5x_1 + x_2 = 6 \implies \begin{cases} 5x_1 + x_2 \geq 6 \\ 5x_1 + x_2 \leq 6 \end{cases} \implies \begin{cases} 5x_1 + x_2 \geq 6 \\ -5x_1 - x_2 \geq -6 \end{cases}$$

*So our problem, in the form of the primal in the symmetric primal/dual pair, is:*

$$\begin{aligned} &\text{Minimize } 20x_1 + 10x_2 \\ &\text{s.t.} \quad 5x_1 + x_2 \geq 6 \\ &\quad \quad -5x_1 - x_2 \geq -6 \\ &\quad \quad 2x_1 + 2x_2 \geq 8 \\ &\quad \quad x_1 \geq 0, x_2 \geq 0 \end{aligned}$$

*We can now write its DUAL problem:*

*(For reasons to be apparent, we choose to name our dual variables not  $y_1, y_2$ , and  $y_3$  but  $y_1', y_1''$ , and  $y_2$ .)*

$$\begin{array}{ll} \mathbf{P:} & \text{Minimize } 20x_1 + 10x_2 \\ & \text{s.t.} \quad 5x_1 + x_2 \geq 6 \\ & \quad \quad -5x_1 - x_2 \geq -6 \\ & \quad \quad 2x_1 + 2x_2 \geq 8 \\ & \quad \quad x_1 \geq 0, x_2 \geq 0 \end{array} \qquad \begin{array}{ll} \mathbf{D:} & \text{Max } 6y_1' - 6y_1'' + 8y_2 \\ & \text{s.t.} \quad 5y_1' - 5y_1'' + 2y_2 \leq 20 \\ & \quad \quad y_1' - y_1'' + 2y_2 \leq 10 \\ & \quad \quad y_1' \geq 0, y_1'' \geq 0, y_2 \geq 0 \end{array}$$

*Notice that the pair of dual variables  $y_1'$  and  $y_1''$  always appear with opposite signs:*

$$\begin{array}{ll}
 \text{Max } 6y_1' - 6y_1'' + 8y_2 & \text{Max } 6(y_1' - y_1'') + 8y_2 \\
 \text{s.t.} & \text{s.t.} \\
 5y_1' - 5y_1'' + 2y_2 \leq 20 & \Rightarrow \quad 5(y_1' - y_1'') + 2y_2 \leq 20 \\
 y_1' - y_1'' + 2y_2 \leq 10 & (y_1' - y_1'') + 2y_2 \leq 10 \\
 y_1' \geq 0, y_1'' \geq 0, y_2 \geq 0 & y_1' \geq 0, y_1'' \geq 0, y_2 \geq 0
 \end{array}$$

*It is instructive now to make the change of variable:  $y_1 = y_1' - y_1''$*

*Letting  $y_1 = y_1' - y_1''$ ,*

$$\begin{array}{ll}
 \text{Max } 6(y_1' - y_1'') + 8y_2 & \text{Maximize } 6y_1 + 8y_2 \\
 \text{s.t.} & \text{subject to} \\
 5(y_1' - y_1'') + 2y_2 \leq 20 & \Rightarrow \quad 5y_1 + 2y_2 \leq 20 \\
 (y_1' - y_1'') + 2y_2 \leq 10 & y_1 + 2y_2 \leq 10 \\
 y_1' \geq 0, y_1'' \geq 0, y_2 \geq 0 & y_2 \geq 0
 \end{array}$$

*(We cannot include a constraint on the sign of  $y_1$ , since it is the difference of two variables.)*

*This is the same as the dual in the symmetric primal/dual pair, except for the missing nonnegativity restriction!*

**We next show:**

**The dual of the DUAL problem  
is the PRIMAL problem!**

**Problem (P):**

Minimize  $c^t x$   
subject to:

$$\begin{aligned} Ax &\geq b \\ x &\geq 0 \end{aligned}$$

**Problem (D):**


Maximize  $b^t y$   
subject to:

$$\begin{aligned} A^t y &\leq c \\ y &\geq 0 \end{aligned}$$

How do we write the DUAL of problem (D) above? First we must write it as a minimization problem with  $\geq$  constraints.

**Problem (D):**Maximize  $b^t y$   
subject to:

$$\begin{aligned} A^t y &\leq c \\ y &\geq 0 \end{aligned}$$

**(equivalent)****Problem (D'):**-Min  $(-b^t)y$   
subject to

$$\begin{aligned} (-A^t)y &\geq -c \\ y &\geq 0 \end{aligned}$$

**MINIMIZING the NEGATIVE of a function yields the same solution (except for sign) as MAXIMIZING the function.**  
**NEGATING both sides of a  $\leq$  constraint produces a  $\geq$  constraint.**

Primal:Minimize  $c^t x$   
subject to:  

$$\begin{aligned} Ax &\geq b \\ x &\geq 0 \end{aligned}$$
Dual:Maximize  $b^t y$   
subject to:  

$$\begin{aligned} A^t y &\leq c \\ y &\geq 0 \end{aligned}$$
**Problem (D'):**-Min  $(-b^t)y$   
subject to  

$$\begin{aligned} (-A^t)y &\geq -c \\ y &\geq 0 \end{aligned}$$
**Problem (DD'):**-Max  $(-c^t)u$   
subject to  

$$\begin{aligned} (-A^t)^t u &\leq (-b^t)^t \\ u &\geq 0 \end{aligned}$$

**Problem (DD')**:

$$\begin{array}{ccc}
 \text{-Max } (-c^t)u & \textit{is equivalent to:} & \text{Min } c^t u \\
 \text{subject to} & \Rightarrow & \text{subject to} \\
 (-A^t)^t u \leq (-b^t)^t & & Au \geq b \\
 u \geq 0 & & u \geq 0
 \end{array}$$

*which is the same as the original PRIMAL problem (P), except for the name of the variables (u instead of x), which is arbitrary!*

**The dual of the DUAL problem  
is the PRIMAL problem!**

*So, given a primal/dual pair of LP problems, it is arbitrary which is referred to as the primal, and which is referred to as the dual.*



<p><b>Writing the dual of a general Primal LP</b></p>
---

The dual of an LP may be found by first rewriting the LP in the form of one of the LPs in the symmetric Primal/Dual pair.

On the other hand, the dual can be written directly for any LP using the following relationships.



Maximize	← →	Minimize
Type of constraint $i$ $\leq$ $=$ $\geq$	← →	Sign of variable $i$ nonnegative unrestricted nonpositive
Sign of variable $j$ nonnegative unrestricted nonpositive	← →	Type of constraint $j$ $\geq$ $=$ $\leq$

**Example***We want to write the dual of the LP:*

$$\text{Minimize } 6X_1 + 3X_2 + 5X_4$$

s.t.

$$X_1 - 2X_2 + 4X_3 \leq 20$$

$$2X_1 + X_2 - X_4 = 30$$

$$5X_2 + X_3 + X_4 \geq 50$$

$$X_1 \geq 0, X_2 \text{ urs}, X_3 \geq 0, X_4 \leq 0$$

**We immediately notice that dual will be MAX  
and size of problem will be 4x3**

$$\text{Minimize } 6X_1 + 3X_2 + 5X_4$$

s.t.

$$X_1 - 2X_2 + 4X_3 \leq 20$$

$$2X_1 + X_2 - X_4 = 30$$

$$5X_2 + X_3 + X_4 \geq 50$$

$$X_1 \geq 0, X_2 \text{ urs}, X_3 \geq 0, X_4 \leq 0$$

*Dual will be a  
maximization*

*# dual variables  
=3*

*#dual constraints  
=4*

Primal

Dual

Minimize $6X_1 + 3X_2 + 5X_4$	Maximize $\dots Y_1 + \dots Y_2 + \dots Y_3$
s.t. $X_1 - 2X_2 + 4X_3 \leq 20$ $2X_1 + X_2 - X_4 = 30$ $5X_2 + X_3 + X_4 \geq 50$	s.t. $\dots Y_1 + \dots Y_2 + \dots Y_3 \dots$ $\dots Y_1 + \dots Y_2 + \dots Y_3 \dots$ $\dots Y_1 + \dots Y_2 + \dots Y_3 \dots$ $\dots Y_1 + \dots Y_2 + \dots Y_3 \dots$
$X_1 \geq 0, X_2 \text{ urs}, X_3 \geq 0, X_4 \leq 0$	$Y_1 \dots, Y_2 \dots, Y_3 \dots$

*Transposing coefficients and right-hand-sides:*

Primal

Dual

Minimize $6X_1 + 3X_2 + 5X_4$	Maximize $20Y_1 + 30Y_2 + 50Y_3$
s.t. $X_1 - 2X_2 + 4X_3 \leq 20$ $2X_1 + X_2 - X_4 = 30$ $5X_2 + X_3 + X_4 \geq 50$	s.t. $1 Y_1 + 2 Y_2 + 0 Y_3 \dots 6$ $-2 Y_1 + 1 Y_2 + 5 Y_3 \dots 3$ $4 Y_1 + 0 Y_2 + 1 Y_3 \dots 0$ $0 Y_1 - 1 Y_2 + 1 Y_3 \dots 5$
$X_1 \geq 0, X_2 \text{ urs}, X_3 \geq 0, X_4 \leq 0$	$Y_1 \dots, Y_2 \dots, Y_3 \dots$

*Determining sign restrictions of dual variables*

Primal

Dual

Minimize $6X_1 + 3X_2 + X_4$
s.t. $X_1 - 2X_2 + 4X_3 \leq 20$ $2X_1 + X_2 - X_4 = 30$ $5X_2 + X_3 + X_4 \geq 50$
$X_1 \geq 0, X_2$ urs, $X_3 \geq 0, X_4 \leq 0$

Maximize $20Y_1 + 30Y_2 + 50Y_3$
s.t. $1 Y_1 + 2 Y_2 + 0 Y_3 \dots\dots 6$ $-2 Y_1 + 1 Y_2 + 5 Y_3 \dots\dots 3$ $4 Y_1 + 0 Y_2 + 1 Y_3 \dots\dots 0$ $0 Y_1 - 1 Y_2 + 1 Y_3 \dots\dots 5$
$Y_1 \leq 0, Y_2$ urs, $Y_3 \geq 0$

Min	Max
$\geq$	nonnegative
$=$	urs
$\leq$	nonpositive

*Determining form of dual constraints:*

Primal

Dual

Minimize $6X_1 + 3X_2 + X_4$
s.t. $X_1 - 2X_2 + 4X_3 \leq 20$ $2X_1 + X_2 - X_4 = 30$ $5X_2 + X_3 + X_4 \geq 50$
$X_1 \geq 0, X_2$ urs, $X_3 \geq 0, X_4 \leq 0$

Maximize $20Y_1 + 30Y_2 + 50Y_3$
s.t. $1 Y_1 + 2 Y_2 + 0 Y_3 \leq 6$ $-2 Y_1 + 1 Y_2 + 5 Y_3 = 3$ $4 Y_1 + 0 Y_2 + 1 Y_3 \leq 0$ $0 Y_1 - 1 Y_2 + 1 Y_3 \geq 5$
$Y_1 \leq 0, Y_2$ urs, $Y_3 \geq 0$

Min	Max
nonnegative	$\leq$
urs	$=$
nonpositive	$\geq$



## The WEAK Duality Theorem:

Consider the symmetric primal/dual pair:

*Primal:*

Minimize  $c^t x$   
subject to:

$$\begin{aligned} Ax &\geq b \\ x &\geq 0 \end{aligned}$$

*Dual:*

Maximize  $b^t y$   
subject to:

$$\begin{aligned} A^t y &\leq c \\ y &\geq 0 \end{aligned}$$

Suppose that  $\hat{x}$  is feasible in the primal problem, and  $\hat{y}$  is feasible in the dual. Then  $c^t \hat{x} \geq b^t \hat{y}$



**Proof**

The proof of the Weak Duality Theorem is very simple:

$$A\hat{x} \geq b \quad \& \quad \hat{y} \geq 0 \quad \Longrightarrow \quad \hat{y}^t A \hat{x} \geq \hat{y}^t b$$

Transpose  $A^t \hat{y} \leq c$  to get  $(A^t \hat{y})^t \leq c^t$ , i.e.  $\hat{y}^t A \leq c^t$

$$\text{Then } \hat{y}^t A \leq c^t \quad \& \quad \hat{x} \geq 0 \quad \Longrightarrow \quad \hat{y}^t A \hat{x} \leq c^t \hat{x}$$

Combining these two inequalities gives us

$$c^t \hat{x} \geq \hat{y}^t A \hat{x} \geq \hat{y}^t b \quad \Longrightarrow \quad c^t \hat{x} \geq b^t \hat{y}$$

## Corollaries of the Weak Duality Theorem:

If  $x^*$  and  $y^*$  are optimal solutions of the primal and dual problems, respectively:

- objective value for any primal feasible solution is greater than or equal to  $b^t y^*$
- objective value for any dual feasible solution is less than or equal to  $c^t x^*$



## Corollaries of the Weak Duality Theorem (continued):

- if  $\hat{x}$  and  $\hat{y}$  are feasible in the primal & dual problems, respectively, and if  $c^t \hat{x} = b^t \hat{y}$ ,  
then  $\hat{x} = \text{primal optimum } (x^*)$   
 $\hat{y} = \text{dual optimum } (y^*)$
- if the primal is feasible and unbounded below, then the dual problem must be infeasible!
- if the dual is feasible and unbounded above, then the primal problem must be infeasible!



## Theorem:

If  $B^*$  is an optimal basis of the primal problem (P) then the simplex multiplier vector  $\pi^*$  relative to the basis  $B^*$  is an optimal solution to the dual problem (D).

*(The simplex multiplier vector  $\pi$  may be computed by the formula*

$$\pi^* = c_B^t (A^B)^{-1} \quad .)$$



**Proof**

## Proof:

Let's write the problem (P) with equality constraints, as required by the simplex method:

Primal:

Minimize  $c^t x$   
subject to:

$$\begin{aligned} Ax &\geq b \\ x &\geq 0 \end{aligned}$$



Minimize  $[c \mid 0] \begin{bmatrix} x \\ s \end{bmatrix}$  ← *surplus variable*  
subject to  
 $[A \mid -I] \begin{bmatrix} x \\ s \end{bmatrix} = b$   
 $x \geq 0$   
↑ *identity matrix*

Suppose  $\pi^*$  is the optimal simplex multiplier vector. Then the optimality conditions (for terminating the simplex algorithm) must be satisfied, namely

$$\underbrace{\left( \begin{array}{c} \text{cost of} \\ \text{variable} \end{array} \right)}_{c_j} - \underbrace{\pi^* \left( \begin{array}{c} \text{column of} \\ \text{constraint} \\ \text{coefficients} \end{array} \right)}_{\text{"z}_j"} \geq 0$$

*"reduced cost"*

These conditions must be satisfied for both the original variables ( $x$ ) and the surplus variables ( $s$ ):

$$\left( \begin{array}{c} \text{cost of} \\ \text{variable} \end{array} \right) - \pi^* \left( \begin{array}{c} \text{column of} \\ \text{constraint} \\ \text{coefficients} \end{array} \right) \geq 0$$

Minimize  $[c \mid 0] \begin{bmatrix} x \\ s \end{bmatrix}$        $c^t - \pi^* A \geq 0$ , i.e.  $c^t \geq \pi^* A$   
 subject to  $[A \mid -I] \begin{bmatrix} x \\ s \end{bmatrix} = b$        $0 - \pi^* (-I) \geq 0$ , i.e.  $\underbrace{\pi^*}_{\text{feasibility conditions for the dual}} \geq 0$

And so if  $\pi^*$  is the optimal simplex multiplier,

$$\begin{aligned}\pi^* A &\leq c^t, \text{ i.e. } A^t \pi^* \leq c \\ \pi^* &\geq 0\end{aligned}$$

i.e.,  $\pi^*$  is feasible in the dual problem.

Recall the computation of  $\pi^*$ :  $\pi^* = c_B^t (A^B)^{-1}$

Recall also that  $x_B^* = (A^B)^{-1} b$

$$\begin{aligned}\text{Therefore } c^t x^* &= c_B^t x_B^* = \underbrace{c_B^t (A^B)^{-1}}_{\pi^*} b = \pi^* b \\ &\uparrow \\ &\text{since nonbasic variables} \\ &\text{are zero}\end{aligned}$$

Therefore,  $\pi^*$  is feasible in the dual, and the objective functions of the primal & dual problems evaluated at  $x^*$  and  $\pi^*$ , respectively, are equal.

Hence, by a corollary of the WEAK DUALITY THEOREM,  $x^*$  and  $\pi^*$  must both be optimal in their respective problems!

Q.E.D.

## Example:

P: Maximize  $4X_1 + 5X_2$   
subject to

$$X_1 + X_2 \leq 8$$

$$3X_1 + 2X_2 \leq 18$$

$$2X_1 + 5X_2 \leq 15$$

$$5X_1 - X_2 \leq 10$$

$$X_1 \geq 0, X_2 \geq 0$$

*This problem has 2 variables & 4 inequality constraints, and so its dual will have 4 variables and 2 inequality constraints.*

*The dual problem:*

D: Minimize  $8Y_1 + 18Y_2 + 15Y_3 + 10Y_4$   
subject to:

$$Y_1 + 3Y_2 + 2Y_3 + 5Y_4 \geq 4$$

$$Y_1 + 2Y_2 + 5Y_3 - Y_4 \geq 5$$


$$Y_1 \geq 0, Y_2 \geq 0, Y_3 \geq 0, Y_4 \geq 0$$

*This dual problem has fewer constraints than its primal, and, when solved by the simplex method, usually requires*

- *fewer iterations* (typically 1.5m to 2m iterations)
- *fewer computations per iteration* (especially if using the revised simplex!)

*The optimal simplex tableau for the dual problem is*

<i>basic</i>	-z	$Y_1$	$Y_2$	$Y_3$	$Y_4$	$S_1$	$S_2$	RHS
-z	1	$\frac{96}{27}$	$\frac{181}{27}$	0	0	$\frac{65}{27}$	$\frac{55}{27}$	$-\frac{505}{27}$ ← <i>reduced cost row</i>
$Y_4$	0	$\frac{3}{27}$	$\frac{11}{27}$	0	1	$-\frac{5}{27}$	$\frac{2}{27}$	$\frac{14}{27}$
$Y_3$	0	$\frac{6}{27}$	$\frac{13}{27}$	1	0	$-\frac{1}{27}$	$-\frac{5}{27}$	$\frac{19}{27}$


  
*surplus variables*

*The optimal solution to the dual is*

$$Y_1 = Y_2 = 0, Y_3 = \frac{19}{27}, Y_4 = \frac{14}{27}$$

*What is the optimal solution of the primal problem?*

*The Simplex Multiplier vector for the optimal dual tableau is*

$$\pi = \left[ \frac{65}{27}, \frac{55}{27} \right]$$

*(Why? the reduced cost of the surplus variable  $S_1$  is its cost minus  $\pi$  times the column of coefficients:*

$$\text{reduced cost of } S_1 \text{ is } 0 - \pi \begin{bmatrix} -1 \\ 0 \end{bmatrix} = \pi_1$$

*Likewise, the reduced cost of the surplus variable for row #  $i$  is the Simplex Multiplier for that row.)*

*Therefore, the optimal solution of the original primal problem is*

$$x_1 = \frac{65}{27} \quad x_2 = \frac{55}{27}$$

*Thus, we may choose to solve either the primal or the dual problem, whichever is easier, and obtain the solution to both simultaneously!*

**Note:**  $-\pi_i$  appears as the reduced cost of a slack variable in row  $i$ .

If there is a surplus variable in row  $i$ , its reduced cost is  $0 - (-1)\pi_i = +\pi_i$ .

If constraint  $i$  is an equation without slack or surplus variable, then  $\pi_i$  will NOT appear in the optimal tableau!



## The Fundamental Duality Theorem:

### Problem (P):

Minimize  $c^t x$   
subject to:

$$\begin{aligned} Ax &\geq b \\ x &\geq 0 \end{aligned}$$

### Problem (D):

Maximize  $b^t y$   
subject to:

$$\begin{aligned} A^t y &\leq c \\ y &\geq 0 \end{aligned}$$

- If both problems (P) & (D) are feasible, then both have an optimal solution and their optimal values are equal, i.e.,  
 $c^t x^* = b^t y^*$
- If one of the problems [either (P) or (D)] has an unbounded objective, then the other problem is infeasible.
- If only one of the problems is feasible, then its objective must be unbounded over the feasible region.



Proof

*Note that it is possible that BOTH primal and dual problems are infeasible.*

**Example:***Primal*Minimize  $20x_1 + 10x_2$ 

subject to:

$$5x_1 + x_2 \geq 6$$

$$2x_1 + 2x_2 \geq 8$$

$$x_1 \geq 0, x_2 \geq 0$$

*Dual*Maximize  $6y_1 + 8y_2$ 

subject to:

$$5y_1 + 2y_2 \leq 20$$

$$y_1 + 2y_2 \leq 10$$

$$y_1 \geq 0, y_2 \geq 0$$

*The primal system has 6 basic solutions, of which 3 are feasible:*


*Primal*Minimize  $20x_1 + 10x_2$ 

subject to:

$$5x_1 + x_2 \geq 6$$

$$2x_1 + 2x_2 \geq 8$$

$$x_1 \geq 0, x_2 \geq 0$$



*The dual system also has six basic solutions, 4 of them feasible:*

Dual


Maximize  $6y_1 + 8y_2$

subject to:

$$5y_1 + 2y_2 \leq 20$$

$$y_1 + 2y_2 \leq 10$$

$$y_1 \geq 0, y_2 \geq 0$$




extreme pt. #	<u>PRIMAL</u>				feasible?	obj.	feasible?	<u>DUAL</u>			
	$x_1$	$x_2$	$x_3$	$x_4$				$y_1$	$y_2$	$y_3$	$y_4$
<b>VI</b>	4	0	14	0	✓	80		0	10	0	-10
<b>II</b>	0	6	0	4	✓	60		10	0	-30	0
<b>I</b>	.5	3.5	0	0	✓	45	✓	2.5	3.75	0	0
<b>IV</b>	0	4	-2	0		40	✓	0	5	10	0
<b>III</b>	1.2	0	0	-5		24	✓	4	0	0	6
<b>VI</b>	0	0	-6	-8		0	✓	0	0	20	10

← optimal

## Example: Unbounded Primal Problem

Minimize  $2X_1 - 6X_2$   
 subject to  
 $X_1 + X_2 \geq 2$   
 $X_1 - X_2 \geq -1$   
 $X_1 \geq 0, X_2 \geq 0$


*The objective  $\rightarrow -\infty$   
 if we travel along the  
 edge of the feasible region  
 to the upper right!*



*The dual of this unbounded primal problem:*


Maximize  $2Y_1 - Y_2$   
 subject to  
 $Y_1 + Y_2 \leq 2$   
 $Y_1 - Y_2 \leq -6$   
 $Y_1 \geq 0, Y_2 \geq 0$

*(Infeasible!)*




## Example: Infeasible Primal Problem

$$\begin{aligned} &\text{Minimize } X_1 + X_2 \\ &\text{subject to} \\ &\quad X_1 - X_2 \geq 1 \\ &\quad -X_1 + X_2 \geq 1 \\ &\quad X_1 \geq 0, X_2 \geq 0 \end{aligned}$$



*Dual of the infeasible primal problem:*

$$\begin{aligned} &\text{Maximize } Y_1 + Y_2 \\ &\text{subject to} \\ &\quad Y_1 - Y_2 \leq 1 \\ &\quad -Y_1 + Y_2 \leq 1 \\ &\quad Y_1 \geq 0, Y_2 \geq 0 \end{aligned}$$



*Objective is unbounded as we move to the upper right!*

*Recall that it is possible that BOTH primal and dual problems are infeasible!*

*Can you alter the preceding (infeasible) primal problem so that the dual problem becomes infeasible (while the primal problem remains infeasible)?*

Hint: Leave the primal constraints unchanged. Can you then change the dual RHS (=primal objective coefficients) so that the dual becomes infeasible?



## **An economic interpretation of the LP dual problem:**

Consider the DIET PROBLEM:

A housewife has to find a minimum-cost diet for her family by selecting from among 5 foods, subject to the constraints that the diet will provide at least 21 units of vitamin A and 12 units of vitamin B per person per day:

Food:	1	2	3	4	5
Vit. A content	1	0	1	1	2 units/oz.
Vit. B content	0	1	2	1	1 units/oz.
Cost	20	20	31	11	12 ¢/oz.



***The housewife's LP model:***

$$\begin{array}{l}
 \text{Minimize } 20x_1 + 20x_2 + 31x_3 + 11x_4 + 12x_5 \quad \leftarrow \text{cost per person per day} \\
 \text{subject to} \\
 \quad x_1 \quad \quad + x_3 \quad + x_4 \quad + 2x_5 \geq 21 \quad \leftarrow \text{Vit. A reqmt.} \\
 \quad \quad x_2 \quad + 2x_3 \quad + x_4 \quad + x_5 \geq 12 \quad \leftarrow \text{Vit. B reqmt.} \\
 \quad \quad \quad x_1, \dots, x_5 \geq 0
 \end{array}$$

where  $x_j$  = quantity of food #j (oz./day) per person

*(She is ignoring requirements for all other nutrients, and consideration of palatability, etc.)*

***The Pill Salesman's Problem:***

Consider a door-to-door salesman of vitamin pills. He has a supply of vitamin A pills (1 unit each) and vitamin B pills (also 1 unit each).

He visits the housewife and suggests that she buy pills from him to feed her family, rather than the foods #1 through #5.

In order to be competitive with the grocery, she must be able to feed her family pills for a cost no more than that of her least-cost meal. *(We ignore the value of her labor!)*

***The Pill Salesman's LP problem:***

Choose prices of the pills:

$\pi_A$  = price per unit of vitamin A pill

$\pi_B$  = price per unit of vitamin B pill

so as to Maximize  $21\pi_A + 12\pi_B$   $\leftarrow$  revenue (\$/day/person)  
subject to

$$\left. \begin{array}{rcl} \pi_A & & \leq 20 \\ & \pi_B & \leq 20 \\ \pi_A + 2\pi_B & \leq & 31 \\ \pi_A + \pi_B & \leq & 11 \\ 2\pi_A + \pi_B & \leq & 12 \\ \pi_A \geq 0, \pi_B \geq 0 \end{array} \right\} \begin{array}{l} \text{the pill-equivalent} \\ \text{of each food must} \\ \text{cost no more than} \\ \text{the food itself} \end{array}$$

*But this is the DUAL of the housewife's LP problem:*

Minimize  $20x_1 + 20x_2 + 31x_3 + 11x_4 + 12x_5$   
subject to

$$\begin{array}{rcl} x_1 & + & x_3 + x_4 + 2x_5 \geq 21 \\ & x_2 & + 2x_3 + x_4 + x_5 \geq 12 \\ & & x_1, \dots, x_5 \geq 0 \end{array}$$

*The Fundamental Duality Theorem tells us that (if both problems are feasible & bounded) the two LP problems have the same optimal values! That is, the housewife would be indifferent between preparing the meals & serving the pills.*



## The FARKAS Lemma:

The following statements are equivalent:

- (i) if  $y^t A \leq 0$  for some  $y$ , then  $y^t b \leq 0$
- &
- (ii) the system  $Ax=b, x \geq 0$  is feasible

*(This "Lemma" is of great theoretical importance in optimization, and is used in the proof of the Kuhn-Tucker optimality conditions in nonlinear programming.)*



**Proof**

### *Proof of the FARKAS Lemma:*

Consider the following primal/dual pair of LPs:

$$\begin{aligned} \text{(P): Minimize } & 0x \\ \text{s.t. } & Ax = b \\ & x \geq 0 \end{aligned}$$

$$\begin{aligned} \text{(D): Maximize } & b^t y \\ \text{s.t. } & A^t y \leq 0 \end{aligned}$$

*or, equivalently,*

$$\begin{aligned} \text{Maximize } & y^t b \\ \text{s.t. } & y^t A \leq 0 \end{aligned}$$

*(These LP problems have interesting characteristics:*

- *every feasible solution to (P) is optimal*
- *the value  $y=0$  is feasible in problem (D)*

*First we will prove that if statement (i) is true, i.e.*  
 if  $y^t A \leq 0$  for some  $y$ , then  $y^t b \leq 0$   
*then statement (ii) must also be true, i.e.,*  
 the system  $Ax=b, x \geq 0$  is feasible

If statement (i) is true, then since  $y=0$  is feasible in (D) with objective value 0, it must be optimal for (D) [since (i) says that every feasible solution of (D) has objective value no greater than zero].

The Fundamental Duality Theorem then implies that problem (P) is feasible, which is simply statement (ii) above.

*We next want to prove that if statement (ii) is true, i.e.,*  
 the system  $Ax=b, x \geq 0$  is feasible  
*then statement (i) must also be true, i.e.,*  
 if  $y^t A \leq 0$  for some  $y$ , then  $y^t b \leq 0$

Suppose that  $Ax=b$  for some  $x \geq 0$ , and  $y^t A \leq 0$ . (*We need to show that  $y^t b \leq 0$  .*)

But  $y^t A \leq 0$  &  $x \geq 0$  together imply that  $y^t A x \leq 0$ , and since  $Ax=b$ , that  $y^t b \leq 0$ . That is, statement (i) is true.

## Complementary Slackness

**Theorem:** Suppose that  $\hat{x}$  and  $\hat{y}$  are feasible solutions in the primal & dual problems, respectively:

*Primal:*

Minimize  $c^t x$   
 subject to:  
 $Ax \geq b$   
 $x \geq 0$

*Dual:*

Maximize  $b^t y$   
 subject to:  
 $A^t y \leq c$   
 $y \geq 0$



*(Complementary Slackness, cont'd.)*

Then  $\hat{x}$  and  $\hat{y}$  are each optimal in their respective problems *if and only if:*

- whenever a constraint of one problem is slack, then the corresponding variable of the other problem is zero
- whenever a variable of one problem is positive, then the corresponding constraint of the other problem is tight.



Proof

***Proof of the Complementary Slackness Theorem:***

*First we introduce surplus & slack variables to the primal & dual problems, respectively:*

$$\begin{array}{ll}
 \text{P: Min } c^t x & \text{D: Max } yb \\
 \text{s.t.} & \text{s.t.} \\
 Ax - Iu = b & yA + Iv = c^t \\
 x \geq 0, u \geq 0 & y \geq 0, v \geq 0
 \end{array}$$

*Now suppose that the vector  $[\hat{x}, \hat{u}]$  is feasible in P, and that  $[\hat{y}, \hat{v}]$  is feasible in D.*

*(Proof of Complementary Slackness, cont'd.):*

*Consider the difference:*

$$\begin{aligned}
 c^t \hat{x} - \hat{y} b &= \underbrace{(\hat{y}A + I\hat{v})}_{c^t} \hat{x} - \hat{y} \underbrace{(A\hat{x} - I\hat{u})}_b \\
 &= \hat{y} A \hat{x} + \hat{v} \hat{x} - \hat{y} A \hat{x} + \hat{y} \hat{u} \\
 &= \hat{v} \hat{x} + \hat{y} \hat{u}
 \end{aligned}$$

*(1) Suppose that  $[\hat{x}, \hat{u}]$  and  $[\hat{y}, \hat{v}]$  are both optimal in their respective problems, i.e.,  $c^t \hat{x} = \hat{y} b$ .*

$$\text{Then } \hat{v} \hat{x} + \hat{y} \hat{u} = 0$$

$$\text{That is, } \sum_{j=1}^n \hat{v}_j \hat{x}_j + \sum_{i=1}^m \hat{y}_i \hat{u}_i = 0$$

*(Proof of Complementary Slackness, cont'd.):*

*Since each of the factors in each term  $\hat{v}_j \hat{x}_j$  and  $\hat{y}_i \hat{u}_i$  are nonnegative, each term is nonnegative.*

*And because the sum of all terms is zero, it is clear that each term must be zero, i.e.,*

$$\sum_{j=1}^n \hat{v}_j \hat{x}_j + \sum_{i=1}^m \hat{y}_i \hat{u}_i = 0 \implies \hat{v}_j \hat{x}_j = 0 \quad \& \quad \hat{y}_i \hat{u}_i = 0$$

*for all  $j=1, \dots, n$       for all  $i=1, \dots, m$*

$$\hat{v}_j \hat{x}_j = 0 \implies \text{either } \hat{v}_j = 0 \text{ or } \hat{x}_j = 0$$

$$\hat{y}_i \hat{u}_i = 0 \implies \text{either } \hat{y}_i = 0 \text{ or } \hat{u}_i = 0$$

*(Proof of Complementary Slackness, cont'd.):*

$$\text{But } \hat{v}_j \hat{x}_j = 0 \implies \text{either } \hat{v}_j = 0 \text{ or } \hat{x}_j = 0$$

*i.e., when dual constraint #j is slack, the corresponding primal variable  $\hat{x}_j$  must be zero.*

*and*

*when the primal variable  $\hat{x}_j$  is positive, then the corresponding dual constraint must be tight.*

*(Proof of Complementary Slackness, cont'd.):*

And  $\hat{y}_i \hat{u}_i = 0 \implies$  either  $\hat{y}_i = 0$  or  $\hat{u}_i = 0$

*i.e., when primal constraint #i is slack ( $\hat{u}_i > 0$ ), then the corresponding dual variable ( $\hat{y}_i$ ) must be zero and when a dual variable ( $\hat{y}_i$ ) is positive, the corresponding primal constraint must be tight (so that  $\hat{u}_i = 0$ ).*

*So optimality implies that complementary slackness is satisfied.*

*(Proof of Complementary Slackness, cont'd.):*

*The converse is also true: if complementary slackness is satisfied, then the solutions must be optimal, since*

$$c^t \hat{x} - \hat{y} b = \sum_{j=1}^n \hat{y}_j \hat{x}_j + \sum_{i=1}^m \hat{y}_i \hat{u}_i$$

*and so, if each term is zero, the sum must be zero, i.e.,*

$$c^t \hat{x} - \hat{y} b = 0 \implies c^t \hat{x} = \hat{y} b$$

*which, according to the Weak Duality Theorem, means that  $\hat{x}$  and  $\hat{y}$  must both be optimal in their respective problems.*

Q.E.D.