

Revised Simplex Method for LP

This Hypercard stack was prepared by:
Dennis Bricker,
Dept. of Industrial Engineering,
University of Iowa,
Iowa City, Iowa 52242.
e-mail: dbricker@icaen.uiowa.edu



In the "ordinary" simplex method, the *full* tableau is updated at every iteration, but much of the information in the tableau is not used in each iteration-- e.g., only the substitution rates in the column entering the basis are actually used!

The "revised" simplex method computes *only* those numbers in the tableau that are required at the current iteration.

Advantages:

- savings in storage requirements in memory
- savings in computation
- control of round-off errors

Basis Set

The set of indices, i.e., column numbers, of the basic variables forms the **basis**, denoted **B**

For example,

if X_2 , X_7 , and X_3 are basic variables, then

$$\mathbf{B} = \{2, 7, 3\}$$

©Dennis Bricker, U. of Iowa, 1998

Basic Variables

The basic variables corresponding to the basis $\mathbf{B} = \{3, 6, 1\}$ form a subvector of \mathbf{X} , denoted by

$$\mathbf{X}_B = \{X_3, X_6, X_1\}$$

The subvector of the cost vector \mathbf{C} corresponding to this basis is denoted

$$\mathbf{C}_B = \{C_3, C_6, C_1\}$$

©Dennis Bricker, U. of Iowa, 1998

Basis Matrix

The columns of the constraint coefficient matrix A indexed by the basis B is the **basis matrix** denoted A^B

For example, if $B=\{3,6,1\}$ and the original tableau is

-Z	x_1	x_2	x_3	x_4	x_5	x_6	x_7	b
1	2	3	1	6	0	0	0	0
0	1	1	3	2	-1	0	0	5
0	5	0	2	9	0	1	0	8
0	3	-1	0	5	0	0	1	3

then

$$A^B = \begin{bmatrix} 3 & 0 & 1 \\ 2 & 1 & 5 \\ 0 & 0 & 3 \end{bmatrix}$$

©Dennis Bricker, U. of Iowa, 1998

Basis Inverse

The inverse of the basis matrix will be called the **basis inverse**

For example, if $B=\{3,6,1\}$ and $A^B = \begin{bmatrix} 3 & 0 & 1 \\ 2 & 1 & 5 \\ 0 & 0 & 3 \end{bmatrix}$

then the basis inverse is

$$(A^B)^{-1} = \begin{bmatrix} 1/3 & 0 & -1/9 \\ -2/3 & 1 & -13/9 \\ 0 & 0 & 1/3 \end{bmatrix}$$

©Dennis Bricker, U. of Iowa, 1998

In the simplex method, we pivot in the tableau in order to change the basis:

-Z	X ₁	X ₂	X ₃	X ₄	X ₅	X ₆	X ₇	b
1	2	3	1	6	0	0	0	0
0	1	1	3	2	-1	0	0	5
0	5	0	2	9	0	1	0	8
0	3	-1	0	5	0	0	1	3

original tableau

tableau with basic variables -Z, X₃, X₆, X₁

-Z	X ₁	X ₂	X ₃	X ₄	X ₅	X ₆	X ₇	b
1	0	3.22	0	2.56	0.333	0	-0.556	-3.33
0	0	0.444	1	0.111	-0.333	0	-0.111	1.33
0	0	0.778	0	0.444	0.667	1	-1.44	0.333
0	1	-0.333	0	1.67	0	0	0.333	1

©Dennis Bricker, U. of Iowa, 1998

If we have the basis inverse matrix and the original tableau, we can compute the current tableau:

$$(A^B)^{-1} = \begin{bmatrix} 1/3 & 0 & -1/9 \\ -2/3 & 1 & -13/9 \\ 0 & 0 & 1/3 \end{bmatrix} \quad A = \begin{bmatrix} 1 & 1 & 3 & 2 & -1 & 0 & 0 \\ 5 & 0 & 2 & 9 & 0 & 1 & 0 \\ 3 & -1 & 0 & 5 & 0 & 0 & 1 \end{bmatrix}$$

$$(A^B)^{-1}A = \begin{bmatrix} 0 & 0.444 & 1 & 0.111 & -0.333 & 0 & -0.111 \\ 0 & 0.778 & 0 & 0.444 & 0.667 & 1 & -1.44 \\ 1 & -0.333 & 0 & 1.67 & 0 & 0 & 0.333 \end{bmatrix}$$

which is that section of the current tableau which contained A originally!

©Dennis Bricker, U. of Iowa, 1998

If the variable X_k has been chosen to enter the basis, then the "minimum ratio test" must be performed to choose the pivot row.

The "**substitution rates**" to be used in the denominator of these ratios may therefore be obtained by $(A^B)^{-1}A^k$.

e.g.,

$$\alpha = (A^B)^{-1}A^4 = \begin{bmatrix} 1/3 & 0 & -1/9 \\ -2/3 & 1 & -13/9 \\ 0 & 0 & 1/3 \end{bmatrix} \begin{bmatrix} 2 \\ 9 \\ 5 \end{bmatrix} = \begin{bmatrix} 1/9 \\ 4/9 \\ 5/3 \end{bmatrix}$$

©Dennis Bricker, U. of Iowa, 1998

We can use the basis inverse matrix to compute the values of the basic variables:

$$(A^B)^{-1} = \begin{bmatrix} 1/3 & 0 & -1/9 \\ -2/3 & 1 & -13/9 \\ 0 & 0 & 1/3 \end{bmatrix} \quad b = \begin{bmatrix} 5 \\ 8 \\ 3 \end{bmatrix}$$

$$(A^B)^{-1} b = \begin{bmatrix} 1.33 \\ 0.333 \\ 1 \end{bmatrix} \leftarrow \text{which is the RHS of the current tableau, i.e., the values of the basic variables:}$$

Note the order, which corresponds to $B = \{3, 6, 1\}$

$$\begin{cases} X_3 = 4/3 \\ X_6 = 1/3 \\ X_1 = 1 \end{cases}$$

©Dennis Bricker, U. of Iowa, 1998

The **reduced cost** of variable X_k is

$$\bar{C}_k = C_k - \sum_{i=1}^m C_{B_i} \alpha_i$$

where α_i is the substitution rate of X_k for the basic variable in row i , i.e., of X_{B_i}

That is, if X_k is increased by 1 unit, C_k is added to the cost, while the basic variable in row i is reduced by the amount α_i , thereby saving (if $\alpha_i > 0$) a cost $C_{B_i} \alpha_i$

©Dennis Bricker, U. of Iowa, 1998

Since $\alpha = (A^B)^{-1} A^k$, $\bar{C}_k = C_k - C_B (A^B)^{-1} A^k$

If we define $\pi = C_B (A^B)^{-1}$, then

$$\begin{aligned} \bar{C}_k &= C_k - \pi A^k \\ &= C_k - \sum_{i=1}^m \pi_i A_i^k \end{aligned}$$

©Dennis Bricker, U. of Iowa, 1998

original tableau:

-Z	X ₁	X ₂	X ₃	X ₄	X ₅	X ₆	X ₇	b
1	2	3	1	6	0	0	0	0
0	1	1	3	2	-1	0	0	5
0	5	0	2	9	0	1	0	8
0	3	-1	0	5	0	0	1	3

$$B = \{3, 6, 1\}$$

$$(A^B)^{-1} = \begin{bmatrix} 1/3 & 0 & -1/9 \\ -2/3 & 1 & -13/9 \\ 0 & 0 & 1/3 \end{bmatrix}$$

$$\pi = C_B (A^B)^{-1} = [1, 0, 2] \begin{bmatrix} 1/3 & 0 & -1/9 \\ -2/3 & 1 & -13/9 \\ 0 & 0 & 1/3 \end{bmatrix} = [1/3, 0, 5/9]$$

©Dennis Bricker, U. of Iowa, 1998

-Z	X ₁	X ₂	X ₃	X ₄	X ₅	X ₆	X ₇	b
1	2	3	1	6	0	0	0	0
0	1	1	3	2	-1	0	0	5
0	5	0	2	9	0	1	0	8
0	3	-1	0	5	0	0	1	3

$$B = \{3, 6, 1\}$$

$$\pi = [1/3, 0, 5/9]$$

Suppose we want to "price" the nonbasic variable X₄:

$$\begin{aligned} \bar{C}_4 &= C_4 - \pi A^4 \\ &= 6 - \left(\frac{1}{3} \times 2 + 0 \times 9 + \frac{5}{9} \times 5 \right) = \frac{23}{9} \\ &= 2.555555 \end{aligned}$$

©Dennis Bricker, U. of Iowa, 1998

In the **Revised Simplex Method**, no pivoting is done in the LP tableau.

When an entry of the tableau is needed, it is computed, using the inverse of the basis matrix.

- relative profit (or reduced cost) of X_j is

$$C_j - \pi A^j$$

where

$$\pi = C_B (A^B)^{-1}$$

- substitution rates for the entering variable X_j (used in the minimum ratio test) are $\alpha = (A^B)^{-1} A^j$

©Dennis Bricker, U. of Iowa, 1998

Revised Simplex Method

Step 0: determine an initial feasible basis B

Step 1: compute the simplex multiplier vector

$$\pi = C_B (A^B)^{-1}$$

Step 2: price the nonbasic columns, i.e., compute the reduced cost of each nonbasic variable

Step 3: select the entering variable X_k

Step 4: compute the substitution rates of X_k

$$\alpha = (A^B)^{-1} A^k$$

©Dennis Bricker, U. of Iowa, 1998


Revised Simplex Method

Step 5: determine the variable to leave the basis

$$\min_{\alpha_i > 0} \left\{ \frac{\beta_i}{\alpha_i} \right\} = \frac{\beta_\ell}{\alpha_\ell} \quad \text{where } \beta = X_B = (A^B)^{-1} b$$

Step 6: replace the ℓ^{th} index of B with k, and update the inverse matrix $(A^B)^{-1}$
Return to step 1.

©Dennis Bricker, U. of Iowa, 1998



©Dennis Bricker, U. of Iowa, 1998

-Z	X ₁	X ₂	X ₃	X ₄	X ₅	X ₆	X ₇	b
1	2	3	1	6	0	0	0	0
0	1	1	3	2	-1	0	0	5
0	5	0	2	9	0	1	0	8
0	3	-1	0	5	0	0	1	3

$B = \{3, 6, 1\}$
 $(A^B)^{-1} = \begin{bmatrix} 1/3 & 0 & -1/9 \\ -2/3 & 1 & -13/9 \\ 0 & 0 & 1/3 \end{bmatrix}$
 $\pi = C_B (A^B)^{-1} = [1/3, 0, 5/9]$

Example:

Pricing X_7 : $\bar{C}_7 = C_7 - \pi A^7$

$$= 0 - \left(\frac{1}{3} \times 0 + 0 \times 0 + \frac{5}{9} \times 1 \right) = -5/9$$

So X_7 will enter the basis

©Dennis Bricker, U. of Iowa, 1998

original tableau

-Z	X ₁	X ₂	X ₃	X ₄	X ₅	X ₆	X ₇	b
1	2	3	1	6	0	0	0	0
0	1	1	3	2	-1	0	0	5
0	5	0	2	9	0	1	0	8
0	3	-1	0	5	0	0	1	3

$B = \{3, 6, 1\}$
 $(A^B)^{-1} = \begin{bmatrix} 1/3 & 0 & -1/9 \\ -2/3 & 1 & -13/9 \\ 0 & 0 & 1/3 \end{bmatrix}$

current tableau

-Z	X ₁	X ₂	X ₃	X ₄	X ₅	X ₆	X ₇	b
1	0	0	0	0	0	0	-5/9	
0	0	1	0	0	0	0		
0	0	0	0	0	1	0		
0	1	0	0	0	0	0		

Since the reduced cost of the variable X_7 is negative, we will enter it into the basis.

To choose the pivot row will require the "substitution rates" in the column.

©Dennis Bricker, U. of Iowa, 1998

original tableau

-Z	X ₁	X ₂	X ₃	X ₄	X ₅	X ₆	X ₇	b
1	2	3	1	6	0	0	0	0
0	1	1	3	2	-1	0	0	5
0	5	0	2	9	0	1	0	8
0	3	-1	0	5	0	0	1	3

$$B = \{3, 6, 1\}$$

$$(A^B)^{-1} = \begin{bmatrix} 1/3 & 0 & -1/9 \\ -2/3 & 1 & -13/9 \\ 0 & 0 & 1/3 \end{bmatrix}$$

Calculation of "substitution rates" $\alpha = (A^B)^{-1} A^j$

$$\alpha = \begin{bmatrix} 1/3 & 0 & -1/9 \\ -2/3 & 1 & -13/9 \\ 0 & 0 & 1/3 \end{bmatrix} \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} = \begin{bmatrix} -1/9 \\ -13/9 \\ 1/3 \end{bmatrix}$$

©Dennis Bricker, U. of Iowa, 1998

current tableau

-Z	X ₁	X ₂	X ₃	X ₄	X ₅	X ₆	X ₇	b
1	0	0	0	0	0	-5/9		
0	0	1	0	0	0	-1/9		
0	0	0	0	0	1	-13/9		
0	1	0	0	0	0	1/3		

Next, to perform the "minimum ratio test" for selecting the pivot row, we need the current right-hand-side: $\beta = X_B = (A^B)^{-1} b$

©Dennis Bricker, U. of Iowa, 1998

original tableau

-Z	x_1	x_2	x_3	x_4	x_5	x_6	x_7	b
1	2	3	1	6	0	0	0	0
0	1	1	3	2	-1	0	0	5
0	5	0	2	9	0	1	0	8
0	3	-1	0	5	0	0	1	3

$$B = \{3, 6, 1\}$$

$$(A^B)^{-1} = \begin{bmatrix} 1/3 & 0 & -1/9 \\ -2/3 & 1 & -13/9 \\ 0 & 0 & 1/3 \end{bmatrix}$$

$$\beta = X_B = (A^B)^{-1} b = \begin{bmatrix} 1/3 & 0 & -1/9 \\ -2/3 & 1 & -13/9 \\ 0 & 0 & 1/3 \end{bmatrix} \begin{bmatrix} 5 \\ 8 \\ 3 \end{bmatrix} = \begin{bmatrix} 2/3 \\ 1/3 \\ 1 \end{bmatrix}$$

©Dennis Bricker, U. of Iowa, 1998

current tableau

-Z	x_1	x_2	x_3	x_4	x_5	x_6	x_7	b
1	0	0			0	$-5/9$		
0	0		1		0	$-1/9$		$2/3$
0	0		0		1	$-13/9$		$1/3$
0	1		0		0	$1/3$		1

Now we can select the pivot row, which is the last row (since there is only one positive substitution rate). That is, x_7 enters the basis, replacing x_1 , so that the new basis is $B = \{3, 6, 7\}$

©Dennis Bricker, U. of Iowa, 1998

original tableau

-Z	X ₁	X ₂	X ₃	X ₄	X ₅	X ₆	X ₇	b
1	2	3	1	6	0	0	0	0
0	1	1	3	2	-1	0	0	5
0	5	0	2	9	0	1	0	8
0	3	-1	0	5	0	0	1	3

$$B = \{3, 6, 7\}$$

$$A^B = \begin{bmatrix} 3 & 0 & 0 \\ 2 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

Rather than to compute the basis inverse matrix "from scratch", it is possible to update the old basis inverse matrix:

$$\left[\begin{array}{ccc|c} 1/3 & 0 & -1/9 & -1/9 \\ -2/3 & 1 & -13/9 & -13/9 \\ 0 & 0 & 1/3 & 1/3 \end{array} \right] \longrightarrow \left[\begin{array}{ccc|c} 1/3 & 0 & 0 & 0 \\ -2/3 & 1 & 0 & 0 \\ 0 & 0 & 1 & 1 \end{array} \right]$$

old inverse
pivot column
new inverse

©Dennis Bricker, U. of Iowa, 1998

We are then ready to begin another iteration of the revised simplex method.

$$\pi = \left[-1/3, 0, 0 \right]$$

$$\bar{C}_2 = 3 - (\pi_1 - \pi_3) = 8/3$$

$$\bar{C}_4 = 6 - (2\pi_1 + 9\pi_2 + 5\pi_3) = 16/3$$

$$\bar{C}_5 = 0 - (-\pi_1) = 1/3$$

Since the reduced costs of the nonbasic variables are all positive, the current basic solution is optimal!

Advantages of Revised Simplex Method

- Because we pivot only in the basis inverse matrix, each pivot requires updating only $m \times m$ numbers, not $(m+1)(n+1)$ (where $m = \# \text{rows}$, $n = \# \text{columns}$)

For example, if $m=100$ and $n=1000$ (not unusually big in practice), the RSM updates only 10,000 numbers in the basis inverse, while the ordinary simplex method updates 100,000 numbers in the tableau.

©Dennis Bricker, U. of Iowa, 1998

Advantages of Revised Simplex Method

- Most real-world LP problems have an original tableau which is very sparse, e.g., only 1% to 5% of the numbers are nonzero. This allows efficiency of storage in memory. After several pivots in the tableau, however, the tableau usually loses most of this sparsity! RSM, using the original tableau, can still take advantage of sparsity.

©Dennis Bricker, U. of Iowa, 1998

Advantages of Revised Simplex Method

- Because computation is done using only the basis inverse and the original data, the build-up of round-off errors in the computation is largely avoided.

(LP software will periodically re-invert the basis inverse "from scratch" to avoid this buildup of round-off errors.)

©Dennis Bricker, U. of Iowa, 1998

The Revised Simplex Method (or some variation or extension of it) is used in virtually all commonly-available software for linear programming.

Other common extensions:

- *storage of the basis inverse in "product-form" to save both storage space and computation*
- *factoring the basis inverse into triangular factors to make computation of substitution rates and simplex multipliers more efficient.*
- *simple and general upper bounds on variables*

©Dennis Bricker, U. of Iowa, 1998

Exercise

original tableau

-Z	X ₁	X ₂	X ₃	X ₄	X ₅	X ₆	RHS
1	1	1	-4	0	0	0	0
0	1	1	2	1	0	0	9
0	1	1	-1	0	1	0	2
0	-1	1	1	0	0	1	4

-Z	X ₁	X ₂	X ₃	X ₄	X ₅	X ₆	RHS
1							
0				1/3	0	-2/3	
0				0	1	1	
0				1/3	0	1/3	

current tableau

Complete the missing elements!

Solution

©Dennis Bricker, U. of Iowa, 1998

Solution:

original tableau

-Z	X ₁	X ₂	X ₃	X ₄	X ₅	X ₆	RHS
1	1	1	-4	0	0	0	0
0	1	1	2	1	0	0	9
0	1	1	-1	0	1	0	2
0	-1	1	1	0	0	1	4

-Z	X ₁	X ₂	X ₃	X ₄	X ₅	X ₆	RHS
1	0	4	0	1	0	2	17
0	1	-1/3	0	1/3	0	-2/3	1/3
0	0	2	0	0	1	1	6
0	0	2/3	1	1/3	0	1/3	13/3

current tableau

©Dennis Bricker, U. of Iowa, 1998

Exercise

original tableau

-Z	X ₁	X ₂	X ₃	X ₄	X ₅	RHS
1	a	-1	2	0	0	0
0	b	c	d	1	0	6
0	-1	3	e	0	1	1

current tableau

-Z	X ₁	X ₂	X ₃	X ₄	X ₅	RHS
1	0	-7	j	k	m	-9
0	g	2	-1	1/2	0	f
0	h	i	1	1/2	1	4

Find the values of "a" through "m"



Solution

©Dennis Bricker, U. of Iowa, 1998

Solution

- a= 3
- b= 2
- c= 4
- d= -2
- e= 2
- f= 3
- g= 1
- h= 0
- i= 5
- j= 5
- k= -3/2
- m= 0

original tableau

-Z	X ₁	X ₂	X ₃	X ₄	X ₅	RHS
1	a	-1	2	0	0	0
0	b	c	d	1	0	6
0	-1	3	e	0	1	1

current tableau

-Z	X ₁	X ₂	X ₃	X ₄	X ₅	RHS
1	0	-7	j	k	m	-9
0	g	2	-1	1/2	0	f
0	h	i	1	1/2	1	4



©Dennis Bricker, U. of Iowa, 1998