

## Variations of the Simplex Tableau

**Summary:** The simplex tableau as shown in various textbooks and these notes may differ, and cause you some confusion.

© Dennis Bricker  
Dept of Mechanical & Industrial Engineering  
The University of Iowa

Let's assume the LP model

$$\begin{aligned} \text{Minimize } z &= \sum_{j=1}^n c_j x_j \\ \text{s.t. } \sum_{j=1}^n a_{ij} x_j &= b_i, \quad \forall i = 1, \dots, m \\ x_j &\geq 0, \quad j = 1, \dots, n \end{aligned}$$

We can write the objective equation in either of two ways:

$$z - \sum_{j=1}^n c_j x_j = 0 \quad \text{or} \quad -z + \sum_{j=1}^n c_j x_j = 0$$

The objective row can be written either as the **first** or **last row** of the tableau. In my notes, it will appear as the first row. In my notes, I will use “-z” as the basic variable in the objective row.

For example, the tableau might appear as

-z	X <sub>1</sub>	X <sub>2</sub>	X <sub>3</sub>	X <sub>4</sub>	X <sub>5</sub>	X <sub>6</sub>	RHS
1	2	0	-3	0	-4	0	-22
0	-1	0	0	1	3	0	5
0	4	1	2	0	0	0	2
0	2	0	1	0	-2	1	3

where X<sub>2</sub>, X<sub>4</sub>, and X<sub>6</sub> (along with -z) are basic.

In this tableau, the values in the objective row are the **reduced costs**. Thus,

- increasing X<sub>1</sub> will **increase** the objective at the rate of 2 units of cost per unit of X<sub>1</sub>,
- increasing X<sub>3</sub> will **decrease** the objective at the rate of 3 units of cost per unit of X<sub>3</sub>.

If **minimizing**, then any column with **negative** value in objective row can be chosen as the pivot column.

Since the first column never changes, some textbooks don't explicitly include it in the tableau:

-z	X <sub>1</sub>	X <sub>2</sub>	X <sub>3</sub>	X <sub>4</sub>	X <sub>5</sub>	X <sub>6</sub>	RHS
1	2	0	-3	0	-4	0	-22
0	-1	0	0	1	3	0	5
0	4	1	2	0	0	0	2
0	2	0	1	0	-2	1	3

i.e.,

X <sub>1</sub>	X <sub>2</sub>	X <sub>3</sub>	X <sub>4</sub>	X <sub>5</sub>	X <sub>6</sub>	RHS
2	0	-3	0	-4	0	-22
-1	0	0	1	3	0	5
4	1	2	0	0	0	2
2	0	1	0	-2	1	3

In some textbooks,  $z$  (rather than  $-z$ ) is used as the basic variable in the objective row. The earlier tableau would then appear as

$z$	$X_1$	$X_2$	$X_3$	$X_4$	$X_5$	$X_6$	RHS
1	-2	0	3	0	4	0	22
0	-1	0	0	1	3	0	5
0	4	1	2	0	0	0	2
0	2	0	1	0	-2	1	3

If this convention is used, then the signs of all values in the objective row are reversed, and the rule for choosing the variable to enter the basis is changed:

If **minimizing**, then any column with **positive** value in objective row can be chosen as the pivot column.  
 If **maximizing**, then any column with **negative** value in objective row can be chosen as the pivot column.

Knowing the basic variables in each row is enough to know the contents of each row:

$z$	$X_1$	$X_2$	$X_3$	$X_4$	$X_5$	$X_6$	RHS
1	-2	0	3	0	4	0	22
0	-1	0	0	1	3	0	5
0	4	1	2	0	0	0	2
0	2	0	1	0	-2	1	3

Therefore, some textbooks don't explicitly show the columns of the basic variables in the tableau, but indicate the basic variable for each row, to get a more **compact tableau**:

	$X_1$	$X_3$	$X_5$	RHS
$-z$	2	-3	-4	-22
$X_4$	-1	0	3	5
$X_2$	4	2	0	2
$X_6$	2	1	-2	3

*(When using this form of the tableau, the formula for performing a pivot will be quite different, since the pivot column is replaced by the result of performing the pivot on the column which was previously basic in the pivot row.)*