

Summer Weather



Summer days are classified as either sunny or cloudy.

Suppose that the weather on any summer day depends on the weather conditions for the previous **two** days.





©Dennis Bricker, U. of Iowa, 1997

If we define a stochastic process $\{X_n; n=1,2,...\}$

by $X_n = \begin{cases} 0 & \text{if day n is cloudy,} \\ 1 & \text{if day n is sunny,} \end{cases}$

then the stochastic process is NOT a Markov chain, since it is not MEMORYLESS,

that is, the probability that (for example) tomorrow is SUNNY depends not only on today's state, but also yesterday's!

If we wish a Markov chain model, the state of the system must incorporate ALL relevant information required to determine the transition probabilities!

©Dennis Bricker, U. of Iowa, 1997

To be exact, suppose that

- if it was **sunny** today and yesterday, then it will be **sunny** tomorrow with probability 80%
- if it was sunny today but cloudy yesterday, then it will be sunny tomorrow with probability of only 60%
- if it was cloudy today but sunny yesterday, then it will be cloudy tomorrow with probability 60%
- if it was cloudy for the last two days, then it will be cloudy tomorrow with probability 90%




©Dennis Bricker, U. of Iowa, 1997

- There are four possible states:
 - i. (S,S): Sunny both yesterday & today
 - ii. (S,C): Sunny yesterday, cloudy today
 - iii. (C,S): Cloudy yesterday, sunny today
 - iv. (C,C): Cloudy both yesterday & today


From the state (S,S), there will be a transition to either (S,S) or (S,C); from (S,C), there will be a transition to either (C,S) or (C,C), etc. That is, tomorrow, today will be yesterday!



©Dennis Bricker, U. of Iowa, 1997



©Dennis Bricker, U. of Iowa, 1997



©Dennis Bricker, U. of Iowa, 1997

etc. (SS) (SC) (CS)

The transition probability matrix is

$$P = \begin{bmatrix} ss & sc & cs & cc \\ 0.8 & 0.2 & 0 & 0 \\ sc & 0 & 0 & 0.4 & 0.6 \\ 0.6 & 0.4 & 0 & 0 \\ cc & 0 & 0 & 0.1 & 0.9 \end{bmatrix}$$

Suppose that Sunday and Monday are both SUNNY. What is the probability that Friday will be sunny?

P ⁴ =	.4936 .2532	.1424 .0928	.0844 .087	.2796 .567
	.4272	.1248	.0928	.3552
	.1398	.0592	.0945	.7065

Beginning in state 1 (S,S), there is a probability of 0.4936+0.0844 = 0.578 that 4 days hence (*Friday*) the system will be in states 1 or 3, i.e. the day will be sunny.

(Or, equivalently, one could find the probability that on **Saturday** the system will be in states 1 (\$,\$) or 2 (\$,C), in which case **Friday** must have been sunny. Using the fifth power of P, we would

F**riday** must have been sumly. Using the fifth power of P, we would find that this gives 0.44552+0.13248 = 0.578, the same value.)

The steady-state distribution is

 $\pi_1 = 0.2727, \pi_2 = 0.0909, \pi_3 = 0.0909, \pi_4 = 0.5454,$

so that in the long run, the probability that a day is sunny is the probability of being in states (S,S) or (C,S) is

$$\pi_1 + \pi_3 = 0.3636$$
,

i.e. 36.36% of the days will be sunny.



©Dennis Bricker, U. of Iowa, 1997



When constructing a Markov chain model of a system, the state must be defined so as to incorporate ALL information necessary to determine the probability distribution of the transitions!

(That is, the probability distribution may depend ONLY on the current state, and not on the state of the system in any prior stages.)



©Dennis Bricker, U. of Iowa, 1997