

8.39 Two equal length, horizontal pipes, one with a diameter of 1 in., the other with a diameter of 2 in., are made of the same material and carry the same fluid at the same flow rate. Which pipe produces the larger head loss? Justify your answer.

For either pipe $h_L = f \frac{l}{D} \frac{V^2}{2g}$, where $V = Q/A = Q/(\frac{\pi}{4} D^2)$.

Thus,

$$h_L = f \frac{l}{D} \left[\frac{4Q}{\pi D^2} \right]^2 / 2g = \frac{8}{\pi^2} f \frac{l}{D^5} Q^2 / g$$

or

$$h_L = \left[\frac{8}{\pi^2} \frac{l Q^2}{g} \right] \frac{f}{D^5} \quad (1)$$

Let $()_1$ and $()_2$ denote the 1 in. and 2 in. diameter pipes, respectively. Thus, with $Q_1 = Q_2$ and $l_1 = l_2$, Eq. (1) gives

$$\frac{h_{L1}}{h_{L2}} = \frac{(f_1 / D_1^5)}{(f_2 / D_2^5)} = \left(\frac{f_1}{f_2} \right) \left(\frac{D_2}{D_1} \right)^5 = \left(\frac{f_1}{f_2} \right) \left(\frac{2 \text{ in.}}{1 \text{ in.}} \right)^5$$

or

$$\frac{h_{L1}}{h_{L2}} = 32 \left(\frac{f_1}{f_2} \right) \quad (2)$$

Although $f_1 \neq f_2$ (because $Re_1 \neq Re_2$ and $\epsilon/D_1 \neq \epsilon/D_2$) the ratio f_1/f_2 would not be significantly different than 1, especially compared to the factor of 32 in Eq. (2). For example, assume $Re_1 = 10,000$ and $\epsilon/D_1 = 0.001$ so that $f_1 = 0.033$ (see Fig. 8.20).

Thus, since

$Re = VD/\nu = (Q/\frac{\pi}{4} D^2) D/\nu = \frac{4Q}{\pi \nu} / D$ it follows that if $Re_1 = 10,000$, then $Re_2 = 5,000$ and $\epsilon/D_2 = 0.0005$ if $\epsilon/D_1 = 0.001$. Hence, $f_2 = 0.037$ so that $h_{L1}/h_{L2} = 32 (0.033/0.037) = 28.5 \gg 1$.

Similar results would be true for other Re , ϵ/D values.

Thus, $h_{L1}/h_{L2} = 32 (f_1/f_2) > 1$, The smaller pipe has the larger head loss.