

Chapter 5: Energy Decay in Isotropic Turbulence

Part 4: Energy Spectrum Equation

In Part 3, an equation for $\mathcal{R}_{ij}(\underline{r}, t)$ was derived by taking the average of $u_i(\underline{x}, t)$ times NS_j at \underline{y} + average of $u_j(\underline{y}, t)$ times NS_i at \underline{x} subject assumptions of homogeneous turbulence:

$$\frac{\partial \mathcal{R}_{ij}}{\partial t}(\underline{r}, t) = \frac{\partial S_{jk,i}}{\partial r_k}(-\underline{r}, t) + \frac{\partial S_{jk,i}}{\partial r_k}(\underline{r}, t) - \frac{1}{\rho} \frac{\partial \mathcal{K}_i}{\partial r_j}(\underline{r}, t) - \frac{1}{\rho} \frac{\partial \mathcal{K}_j}{\partial r_i}(-\underline{r}, t) + 2\nu \frac{\partial^2 \mathcal{R}_{ij}}{\partial r_k^2}(\underline{r}, t)$$

Where:

$$\mathcal{K}_i(\underline{x}, \underline{y}, t) = \overline{u_i(\underline{x}, t)p(\underline{y}, t)}$$

is the two-point pressure-velocity correlation vector.

Recall Fourier transform definitions of the velocity-spectrum $\mathcal{E}_{ij}(\underline{\kappa}, t)$ and $\mathcal{R}_{ij}(\underline{r}, t)$ tensors:

$$\mathcal{E}_{ij}(\underline{\kappa}, t) = \frac{1}{(2\pi)^3} \int_{\mathbb{V}} \mathcal{R}_{ij}(\underline{r}, t) e^{i\underline{\kappa} \cdot \underline{r}} d\underline{r}$$

$$\mathcal{R}_{ij}(\underline{r}, t) = \int_{\mathbb{V}} \mathcal{E}_{ij}(\underline{\kappa}, t) e^{-i\underline{\kappa} \cdot \underline{r}} d\underline{\kappa}$$

$d\underline{\kappa} = d\kappa_1 d\kappa_2 d\kappa_3$

These two equations provide a means of decomposing turbulence correlations into contributions from a continuous range of scales as represented by Fourier components $e^{i\underline{\kappa} \cdot \underline{r}}$.

Fourier transform of the \mathcal{R}_{ij} equation gives,

$$\frac{\partial \mathcal{E}_{ij}}{\partial t}(\underline{\kappa}, t) = T_{ij}(\underline{\kappa}, t) + P_{ij}(\underline{\kappa}, t) - 2\nu\kappa^2 \mathcal{E}_{ij}(\underline{\kappa}, t) \quad (1)$$

Which is a 2nd order tensor equation, where:

$$T_{ij}(\underline{\kappa}, t) = \frac{1}{(2\pi)^3} \int_{\mathbb{V}} S_{ij}(\underline{r}, t) e^{i\underline{\kappa} \cdot \underline{r}} d\underline{r} \quad (2)$$

$$S_{ij}(\underline{r}, t) = \frac{\partial S_{jk,i}}{\partial r_k}(-\underline{r}, t) + \frac{\partial S_{ik,j}}{\partial r_k}(\underline{r}, t) \quad (3)$$

$T_{ij}(\underline{\kappa}, t)$ = rate of transfer of energy (gains or losses) between different scales of turbulent motion due to vortex stretching and re-orientation.

$$P_{ij}(\underline{\kappa}, t) = -\frac{1}{(2\pi)^3} \int_{\mathbb{V}} \left[\frac{1}{\rho} \frac{\partial \mathcal{K}_i}{\partial r_j}(\underline{r}, t) + \frac{1}{\rho} \frac{\partial \mathcal{K}_j}{\partial r_i}(-\underline{r}, t) \right] e^{i\underline{\kappa} \cdot \underline{r}} d\underline{r}$$

is the Fourier transformed pressure velocity term.

$P_{ij}(\underline{\kappa}, t) = 0$ for isotropic turbulence = influence of pressure field on bringing anisotropic turbulence to an isotropic state.

The viscous dissipation term is evaluated as follows:

$$\frac{2\nu}{(2\pi)^3} \int_{\mathbb{V}} \frac{\partial^2 \mathcal{R}_{ij}}{\partial r_k^2}(\underline{r}, t) e^{i\underline{\kappa} \cdot \underline{r}} d\underline{r} = \frac{2\nu}{(2\pi)^3} \int_{\mathbb{V}} i^2 \kappa^2 \mathcal{R}_{ij} e^{i\underline{\kappa} \cdot \underline{r}} d\underline{r} = -2\nu\kappa^2 \mathcal{E}_{ij}(\underline{\kappa}, t)$$

Contracting indices in Eq. (1), the pressure term drops out and integrating over a spherical shell converts \mathcal{E}_{ij} to $E(\kappa, t) = \frac{1}{2} \int_{|\underline{\kappa}|=\kappa} \mathcal{E}_{ii}(\underline{\kappa}, t) d\Omega$ with $K = \int_0^\infty E(\kappa, t) d\kappa$

$$\frac{\partial E}{\partial t}(\kappa, t) = T(\kappa, t) - 2\nu\kappa^2 E(\kappa, t) \quad (4)$$

Which is a scalar equation, where:

$$T(\kappa, t) = \frac{1}{2} \int_{|\underline{\kappa}|=\kappa} T_{ii}(\underline{\kappa}, t) d\Omega = 2\pi\kappa^2 T_{ii}(\kappa, t) \quad (5)$$

$d\Omega$ =elemental
solid angle

is the transfer term and $T_{ii} = \mathcal{F}_k\{S_{ii}(\underline{r}, t)\}$.

Next assume in addition to homogeneous also isotropic turbulence and substitute the isotropic form of $S_{ik,j}$, obtained in Chapter 4 Part 2, into Eq. (3) which gives,

$$S_{ii}(r, t) = u_{rms}^3 \frac{1}{r^2} \frac{d}{dr} \left(r^3 \frac{dk}{dr}(r, t) + 4r^2 k(r, t) \right) \quad (6)$$

Appendix A.1

Where $k(r, t) = S_{111}(r\hat{e}_1)/u_{rms}^3$ is the previously defined (Chapter 4 Part 2) correlation function, not to be confused with wave number κ . Note that $S_{ii} = f(r)$ only in isotropic turbulence.

Using Eq. (6), Eq. (2) becomes (Bernard P4.7),

$$T_{ii}(\kappa, t) = \frac{1}{(2\pi)^2} \int_0^\infty S_{ii}(r, t) r^2 \frac{\sin kr}{kr} dr = f(\kappa) \neq f(\underline{\kappa}) \quad (7)$$

Appendix A.2

Substituting Eq. (7) into Eq. (5) gives,

$$T(\kappa, t) = \frac{1}{2\pi} \int_0^\infty S_{ii}(r, t) \kappa r \sin \kappa r dr \quad (8)$$

Which can be interpreted as the Fourier transform of S_{ii} such that the inverse transform is,

$$S_{ii}(r, t) = 2 \int_0^\infty T(\kappa, t) \frac{\sin \kappa r}{\kappa r} d\kappa \quad (9)$$

Substituting Eq. (6) into Eq. (8) and integrating by parts twice gives

$$T(\kappa, t) = \frac{u_{rms}^3}{\pi} \int_0^\infty \kappa [(3 - \kappa^2 r^2) \sin(\kappa r) - 3\kappa r \cos(\kappa r)] k(r, t) dr \quad (10) \quad \text{Appendix A.3}$$

Which shows that $k(r, t)$ determines the rate of energy transfer between the scales of turbulence, as shown later.

Integrating Eq. (4) between 0 and ∞ and using

$$K(t) = \int_0^\infty E(\kappa, t) d\kappa$$

$$\varepsilon = 2\nu \int_0^\infty \kappa^2 E(\kappa, t) d\kappa$$

Gives

$$\frac{dK(t)}{dt} = \int_0^\infty T(\kappa, t) d\kappa - \varepsilon$$

And in isotropic turbulence

$$\frac{dK}{dt} = -\varepsilon$$

Therefore, in isotropic turbulence

$$\int_0^{\infty} T(\kappa, t) d\kappa = 0$$

i.e., net energy transfer between scales equal zero or in other words gains and losses are conserved.

When $f(r, t)$ is known, $E(\kappa, t)$ can be evaluated using:

$$E(\kappa) = \frac{\overline{u^2}}{\pi} \int_0^{\infty} (3f(r) + rf'(r)) \kappa r \sin(\kappa r) dr \quad (11) \quad \text{Appendix A.4}$$

For the final decay, $f(r, t) = e^{-\frac{r^2}{2\lambda_g^2}}$, such that $E(\kappa)$ for the final decay is given by:

$$E(\kappa, t) = \frac{u_{rms}^2 \lambda_g}{\sqrt{2\pi}} (\kappa \lambda_g)^4 e^{-\frac{1}{2}(\kappa \lambda_g)^2} \quad (12^*) \quad \text{Appendix A.5}$$

Consistent with $E(\kappa) = -4\pi\kappa^4 E_1(\kappa)$, as per Bernard Eq. (4.67).

Substituting Eq. (12) into Eq. (4) yields

$$T(\kappa, t) = E(\kappa, t) \frac{u_{rms}}{\lambda_g} \left((\kappa \lambda_g)^2 - 5 \right)$$

*According to Bernard, this expression is consistent with Eq. (4.67), such that

$$E_1(\kappa) = -\frac{E(\kappa)}{4\pi\kappa^4} = \frac{u_{rms}^2 \lambda_g}{4\pi\sqrt{2\pi}} \lambda_g^4 e^{-\frac{1}{2}(\kappa \lambda_g)^2}$$




Figure 5.8 Energy spectrum budget in final period. —, $\partial E/\partial t$; ---, transfer term; - · -, dissipation. In this illustration $R_\lambda = 10$, with λ denoting λ_g .

In final decay scales for which $\kappa < \sqrt{5}/\lambda_g$ lose energy to those for which $\kappa > \sqrt{5}/\lambda_g$.

$$\kappa_e = 2/\lambda_g \text{ peak } E$$

$$\kappa_d = \sqrt{6}/\lambda_g \text{ peak dissipation}$$

In this example, κ_e and κ_d not well separated due to low R_λ .

No inertial range for above form of $E(\kappa, t)$ equation. As λ_g rises the balance in Fig. 5.8 shifts to smaller wave number, i.e., larger scale \rightarrow higher wave numbers lose all their energy before lower wave numbers.

Hinze pp. 218-220

Consider the integrated form of Eq. (4)

$$\frac{d}{dt} \int_0^\kappa E(\kappa, t) d\kappa = \int_0^\kappa T(\kappa, t) d\kappa - 2\nu \int_0^\kappa \kappa^2 E(\kappa, t) d\kappa \quad (13)$$

If, as previously done, the upper limit of the integral is increased to $\kappa = \infty$

$$\int_0^\infty T(\kappa, t) d\kappa = 0 \quad (14)$$

i.e., net energy transfer between scales is zero or in other words gains and losses are conserved.

An alternative derivation for Eq. (14) can be obtained starting from Eq. (9). Specifying $r = 0$ in Eq. (9) recovers Eq. (14)

$$\int_0^\infty T(\kappa, t) \underbrace{\lim_{r \rightarrow 0} \frac{\sin \kappa r}{\kappa r}}_{\boxed{1}} d\kappa = \frac{1}{2} S_{ii}(0, t)$$

If $S_{ii}(0, t) = 0$.

An alternative form of Eq. (6) for $S_{ii}(r, t)$ is:

$$S_{ii}(r, t) = u_{rms}^3 \left(7 \frac{dk}{dr}(r, t) + r \frac{d^2k}{dr^2}(r, t) + \frac{8}{r} k(r, t) \right) \quad (15) \quad \boxed{\text{Appendix A.6}}$$

Substituting the Taylor expansion of $k(r, t) = \frac{r^3}{3!} k'''(0, t) + \dots$ into Eq. (15) gives

$$S_{ii}(r, t) = \frac{35}{6} r^2 k'''(0, t) \quad \boxed{\text{Appendix A.7}}$$

Which shows that $S_{ii}(r, t)$ behaves like r^2 for small values of r . Consequently,

$$S_{ii}(0, t) = 0$$

And Eq. (14) must hold.

Therefore, for $\kappa = \infty$, Eq. (13) becomes,

$$\frac{d}{dt} \int_0^\infty E(\kappa, t) d\kappa = -2\nu \int_0^\infty \kappa^2 E(\kappa, t) d\kappa$$

$$\frac{dK(t)}{dt} = -\varepsilon$$

The LHS shows the change of total kinetic energy of turbulence and since there are no external energy sources, LHS must equal the dissipation caused by viscous effects.

$T(\kappa, t)$ is the Fourier transform of $S_{ii}(r, t)$, which is related to $k(r, t)$. In the K-H equation, it was previously stated that the term:

$$u_{rms}^3 \left[\frac{\partial k}{\partial r} + \frac{4}{r} k \right]$$

represents inertial processes. However, it can also be interpreted as a “convective” action in the transport of $f(r, t)$, caused by the interaction of eddies of different sizes.

Similarly, the term $\int_0^\kappa T(\kappa, t) d\kappa$ can be interpreted as the interaction of eddies of different wave numbers, transferring energy by inertial effects to or from the eddies in region 0 to κ , which is the reason $T(\kappa, t)$ is referred to as the energy-transfer-spectrum function.

Neglecting the interaction of eddies in Eq. (4) gives

$$\frac{\partial E}{\partial t}(\kappa, t) = -2\nu\kappa^2 E(\kappa, t)$$

And integrating

$$E(\kappa, t) = E(\kappa, t_0) \exp[-2\nu\kappa^2(t - t_0)] \quad (15)$$

Comparing this expression with Eq. (12)

$$E(\kappa, t) = \frac{u_{rms}^2 \lambda_g}{\sqrt{2\pi}} (\kappa \lambda_g)^4 e^{-\frac{1}{2}(\kappa \lambda_g)^2} \quad (12)$$

The two exponential forms are equivalent for $t \gg t_0 \rightarrow t - t_0 \approx t$ since (Hinze p.210)

$$\lambda_f = \sqrt{8\nu t} \Rightarrow \lambda_g = \sqrt{4\nu t}$$

$\lambda_f = \sqrt{2} \lambda_g$

$$\exp[-2\nu\kappa^2 t] = \exp\left[-\frac{1}{2}\kappa^2 \sqrt{4\nu t} \sqrt{4\nu t}\right] = \exp -\frac{1}{2}(\kappa \lambda_g)^2$$

Moreover, assuming IC

$$E(\kappa, t_0) = \frac{u_{rms}^2 \lambda_g}{\sqrt{2\pi}} (\kappa \lambda_g)^4$$

Eq. (12) and Eq. (15) become identical.

This shows that the decrease of kinetic energy with time occurs at a higher rate for large wave number eddies and $E(\kappa, t)$ increases very rapidly, proportional to κ^4 and decreases monotonously to zero as κ increases.

Appendix A

A.1

$$S_{ijl}(\underline{r}) = u_{rms}^3 \left[\left(k - r \frac{dk}{dr} \right) \frac{r_i r_j r_l}{2r^3} - \frac{k}{2} \delta_{ij} \frac{r_l}{r} + \frac{1}{4r} \frac{d(kr^2)}{dr} \left(\delta_{il} \frac{r_j}{r} + \delta_{jl} \frac{r_i}{r} \right) \right]$$

For $j = k$ and $l = i$

$$\begin{aligned} S_{iki}(\underline{r}) &= u_{rms}^3 \left[\left(k - r \frac{dk}{dr} \right) \frac{r_k}{2r} - \frac{k}{2} \delta_{ik} \frac{r_i}{r} + \frac{1}{4r} \frac{d(kr^2)}{dr} \left(\delta_{ii} \frac{r_k}{r} + \delta_{ik} \frac{r_i}{r} \right) \right] \\ &= u_{rms}^3 \left[\left(k - r \frac{dk}{dr} \right) \frac{r_k}{2r} - \frac{k}{2} \frac{r_k}{r} + \frac{1}{4r} \frac{d(kr^2)}{dr} \left(4 \frac{r_k}{r} \right) \right] \\ &= u_{rms}^3 \left[-\frac{dk}{dr} \frac{r_k}{2} + \frac{r_k}{r^2} \frac{d(kr^2)}{dr} \right] \\ &= u_{rms}^3 \left[\frac{r_k}{2} \frac{dk}{dr} + 2k \frac{r_k}{r} \right] \end{aligned}$$

Taking a derivative with respect to r_k yields

$$\frac{\partial S_{iki}}{\partial r_k}(\underline{r}) = u_{rms}^3 \left[\frac{r}{2} k''(r) + \frac{7}{2} k'(r) + \frac{4k(r)}{r} \right] \quad (1A)$$

As shown in Chapter 5 Part 1 Appendix A.2.

Similarly,

$$\frac{\partial S_{iki}}{\partial r_k}(-\underline{r}) = u_{rms}^3 \left[-\frac{r}{2} k''(-r) + \frac{7}{2} k'(-r) - \frac{4k(-r)}{r} \right] \quad (2A)$$

And using the following relations

$$k(r) = -k(-r)$$

$$k'(r) = k'(-r)$$

$$k''(r) = -k''(-r)$$

Into Eq. (2A) gives

$$\frac{\partial S_{iki}}{\partial r_k}(-\underline{r}) = u_{rms}^3 \left[\frac{r}{2} k''(r) + \frac{7}{2} k'(r) + \frac{4k(r)}{r} \right] \quad (3A)$$

Now, defining

$$S_{ii}(\underline{r}, t) = \frac{\partial S_{ik,i}}{\partial r_k}(-\underline{r}, t) + \frac{\partial S_{ik,i}}{\partial r_k}(\underline{r}, t)$$

And using Eqs. (1A) and (3A) yields

$$S_{ii}(\underline{r}, t) = u_{rms}^3 \left[rk''(r) + 7k'(r) + \frac{8k(r)}{r} \right]$$

Multiplying and dividing by r^2 results in

$$S_{ii}(\underline{r}, t) = \frac{1}{r^2} u_{rms}^3 [r^3 k'' + 7r^2 k' + 8kr]$$

Or equivalently

$$S_{ii}(r, t) = u_{rms}^3 \frac{1}{r^2} \frac{d}{dr} (r^3 k' + 4r^2 k)$$

A.2

$$T_{ij}(\underline{\kappa}, t) = \frac{1}{(2\pi)^3} \int_{\mathcal{V}} S_{ij}(\underline{r}, t) e^{i\underline{\kappa} \cdot \underline{r}} d\underline{r} \quad (4A)$$

Converting to spherical coordinates

$$\underline{r} = r(\sin \theta \cos \phi \hat{e}_1 + \sin \theta \sin \phi \hat{e}_2 + \cos \theta \hat{e}_3)$$

Also, for isotropic turbulence

$$S_{ii}(r, t) = u_{rms}^3 \frac{1}{r^2} \frac{d}{dr} \left(r^3 \frac{dk}{dr}(r, t) + 4r^2 k(r, t) \right)$$

Such that S_{ii} is only a function of r , a scalar quantity.

Assume that,

$$\underline{\kappa} = \kappa \hat{e}_3$$

Which is possible due to the isotropy hypothesis, i.e., invariance under rotation and reflection.

Therefore, Eq. (4A) for T_{ii} becomes,

$$\begin{aligned} T_{ii}(\kappa) &= \frac{1}{(2\pi)^3} \int_{\mathcal{V}} S_{ii}(r) e^{i\underline{\kappa} \cdot \underline{r}} d\underline{r} \\ &= \frac{1}{(2\pi)^3} \int_0^\infty dr S_{ii}(r) \int_0^{2\pi} d\phi \int_0^\pi d\theta e^{i\kappa r \cos \theta} r^2 \sin(\theta) \\ &= \frac{1}{(2\pi)^2} \int_0^\infty r^2 dr S_{ii}(r) \int_0^\pi d\theta e^{i\kappa r \cos \theta} \sin(\theta) \end{aligned}$$

Multiply and divide by $-i\kappa r$

$$T_{ii}(\kappa) = -\frac{1}{i\kappa r (2\pi)^2} \int_0^\infty r^2 dr S_{ii}(r) \int_0^\pi -i\kappa r e^{i\kappa r \cos \theta} \sin(\theta) d\theta$$

And use the relation:

$$\frac{d}{d\theta} e^{i\kappa r \cos \theta} = -i\kappa r \sin(\theta) e^{i\kappa r \cos \theta}$$

Gives

$$\begin{aligned}
T_{ii}(\kappa) &= -\frac{1}{i\kappa r(2\pi)^2} \int_0^\infty r^2 dr S_{ii}(r) \int_0^\pi \frac{d}{d\theta} e^{i\kappa r \cos \theta} d\theta \\
&= -\frac{1}{i\kappa r(2\pi)^2} \int_0^\infty r^2 dr S_{ii}(r) [e^{i\kappa r \cos \theta}]_0^\pi \\
&= \frac{2}{\kappa r(2\pi)^2} \int_0^\infty r^2 S_{ii}(r) \underbrace{\frac{e^{i\kappa r} - e^{-i\kappa r}}{2i}}_{\boxed{\sin(\kappa r)}} dr \\
T_{ii}(\kappa) &= \frac{2}{(2\pi)^2} \int_0^\infty r^2 S_{ii}(r) \frac{\sin(\kappa r)}{\kappa r} dr
\end{aligned}$$

A.3

$$T_{ii}(\kappa, t) = \frac{1}{(2\pi)^2} \int_0^\infty S_{ii}(r, t) r^2 \frac{\sin \kappa r}{\kappa r} dr \quad (5A)$$

$$S_{ii}(r, t) = u_{rms}^3 \frac{1}{r^2} \frac{d}{dr} \left(r^3 \frac{dk}{dr}(r, t) + 4r^2 k(r, t) \right) \quad (6A)$$

$$T(\kappa, t) = \frac{1}{2} \int_{|\underline{\kappa}|=\kappa} T_{ii}(\underline{\kappa}, t) d\Omega = 2\pi\kappa^2 T_{ii}(\kappa, t) \quad (7A)$$

$$T(\kappa, t) = \frac{u_{rms}^3}{\pi} \int_0^\infty \kappa [(3 - \kappa^2 r^2) \sin(\kappa r) - 3\kappa r \cos(\kappa r)] k(r, t) dr$$

Substituting Eq. (6A) into (5A) gives

$$\begin{aligned} T_{ii}(\kappa, t) &= \frac{1}{2\pi^2} \int_0^\infty u_{rms}^3 \frac{d}{dr} (r^3 k'(r, t) + 4r^2 k(r, t)) \frac{\sin \kappa r}{\kappa r} dr \\ &= \frac{u_{rms}^3}{2\pi^2} \int_0^\infty \underbrace{\frac{d}{dr} (r^3 k'(r, t) + 4r^2 k(r, t))}_{\boxed{dv}} \underbrace{\frac{\sin \kappa r}{\kappa r}}_{\boxed{u}} dr \end{aligned}$$

Integration by parts

$$\int u dv = uv - \int v du$$

$$\begin{aligned} T_{ii}(\kappa, t) &= \left[\frac{u_{rms}^3}{2\pi^2 \kappa} (r^2 k'(r, t) + 4r k(r, t)) \sin \kappa r \right]_0^\infty \\ &\quad - \frac{u_{rms}^3}{2\pi^2} \int_0^\infty (r^3 k'(r, t) + 4r^2 k(r, t)) \frac{d}{dr} \left(\frac{\sin \kappa r}{\kappa r} \right) dr \quad (8A) \end{aligned}$$

Where:

$$\begin{aligned} & \left[\frac{u_{rms}^3}{2\pi^2\kappa} (r^2 k'(r, t) + 4rk(r, t)) \sin \kappa r \right]_0^\infty \\ &= \frac{u_{rms}^3}{2\pi^2\kappa} \left[(r^2 k'(r, t) + 4rk(r, t)) \sin \kappa r \right]_\infty \\ & \quad - \frac{u_{rms}^3}{2\pi^2\kappa} \left[(0^2 k'(0, t) + 4 \cdot 0 k(0, t)) \sin 0 \right] \end{aligned}$$

And using the fact that at large r , the triple correlation $k(r, t)$ behaves like r^{-4} (see Part 3)

$$\lim_{r \rightarrow \infty} r^2 k'(r, t) = \lim_{r \rightarrow \infty} rk(r, t) = 0$$

Therefore,

$$\left[\frac{u_{rms}^3}{2\pi^2\kappa} (r^2 k'(r, t) + 4rk(r, t)) \sin \kappa r \right]_0^\infty = 0$$

And Eq. (8A) becomes:

$$T_{ii}(\kappa, t) = -\frac{u_{rms}^3}{2\pi^2} \int_0^\infty \left(r^3 k'(r, t) + 4r^2 k(r, t) \right) \frac{d}{dr} \left(\frac{\sin \kappa r}{\kappa r} \right) dr \quad (9A)$$

Evaluating the derivative of $\sin \kappa r / \kappa r$ as

$$\frac{d}{dr} \left(\frac{\sin \kappa r}{\kappa r} \right) = \frac{\cos \kappa r}{r} - \frac{\sin \kappa r}{\kappa r^2}$$

And substituting into Eq. (9A) gives

$$\begin{aligned}
T_{ii}(\kappa, t) &= -\frac{u_{rms}^3}{2\pi^2} \int_0^\infty (r^3 k'(r, t) + 4r^2 k(r, t)) \left[\frac{\cos \kappa r}{r} - \frac{\sin \kappa r}{\kappa r^2} \right] dr \\
&= -\frac{u_{rms}^3}{2\pi^2} \int_0^\infty \left(r^2 k'(r, t) \cos \kappa r + 4rk(r, t) \cos \kappa r - \frac{r}{\kappa} k'(r, t) \sin \kappa r \right. \\
&\quad \left. - \frac{4}{\kappa} k(r, t) \sin \kappa r \right) dr
\end{aligned}$$

Using the Product Rule of derivatives

$$\begin{aligned}
T_{ii}(\kappa, t) &= -\frac{u_{rms}^3}{2\pi^2} \int_0^\infty \left(\frac{d(r^2 k(r, t))}{dr} - 2rk(r, t) \right) \cos \kappa r + 4rk(r, t) \cos \kappa r \\
&\quad - \frac{1}{\kappa} \left(\frac{d(rk(r, t))}{dr} - k(r, t) \right) \sin \kappa r - \frac{4}{\kappa} k(r, t) \sin \kappa r dr \\
&= -\frac{u_{rms}^3}{2\pi^2} \int_0^\infty \left(\frac{d(r^2 k(r, t))}{dr} + 2rk(r, t) \right) \cos \kappa r \\
&\quad - \frac{1}{\kappa} \left(\frac{d(rk(r, t))}{dr} + 3k(r, t) \right) \sin \kappa r dr
\end{aligned}$$

Grouping terms depending on $k(r, t)$ and integrating by parts again

$$\begin{aligned}
T_{ii}(\kappa, t) &= -\frac{u_{rms}^3}{2\pi^2} \left(\int_0^\infty 2rk(r, t) \cos \kappa r - \frac{3k(r, t)}{\kappa} \sin \kappa r dr \right) \\
&\quad - \frac{u_{rms}^3}{(2\pi)^2} \int_0^\infty \underbrace{\left(\frac{d(r^2 k(r, t))}{dr} \right)}_{\boxed{dv}} \underbrace{\cos \kappa r}_{\boxed{u}} dr \\
&\quad + \frac{u_{rms}^3}{\kappa(2\pi)^2} \int_0^\infty \underbrace{\left(\frac{d(rk(r, t))}{dr} \right)}_{\boxed{dv}} \underbrace{\sin \kappa r}_{\boxed{u}} dr \\
&= -\frac{u_{rms}^3}{2\pi^2} \left(\int_0^\infty 2rk(r, t) \cos \kappa r - \frac{3k(r, t)}{\kappa} \sin \kappa r dr \right) \\
&\quad - \frac{u_{rms}^3}{2\pi^2} \left[\cancel{r^2 k(r, t) \cos \kappa r} \Big|_0^\infty + \int_0^\infty \kappa r^2 k(r, t) \sin \kappa r dr \right] \\
&\quad + \frac{u_{rms}^3}{2\pi^2 \kappa} \left[\cancel{rk(r, t) \sin \kappa r} \Big|_0^\infty - \int_0^\infty \kappa rk(r, t) \cos \kappa r dr \right]
\end{aligned}$$

Where the fact that at large r , the triple correlation $k(r, t)$ behaves like r^{-4} was invoked.

$$\begin{aligned}
T_{ii} &= -\frac{u_{rms}^3}{2\pi^2} \int_0^\infty \left[3rk(r, t) \cos \kappa r - \frac{3k(r, t)}{\kappa} \sin \kappa r \right. \\
&\quad \left. + \kappa r^2 k(r, t) \sin \kappa r \right] dr \quad (10A)
\end{aligned}$$

Substituting Eq. (10A) into (7A) gives

$$T(\kappa, t) = \frac{\kappa^2 u_{rms}^3}{\pi} \int_0^\infty \left[-3rk(r, t) \cos \kappa r + \frac{3k(r, t)}{\kappa} \sin \kappa r - \kappa r^2 k(r, t) \sin \kappa r \right] dr$$

$$= \frac{u_{rms}^3}{\pi} \int_0^\infty \left[-3\kappa^2 rk(r, t) \cos \kappa r + 3\kappa k(r, t) \sin \kappa r - \kappa^3 r^2 k(r, t) \sin \kappa r \right] dr$$

$$T(\kappa, t) = \frac{u_{rms}^3}{\pi} \int_0^\infty \kappa \left[(3 - \kappa^2 r^2) \sin(\kappa r) - 3\kappa r \cos(\kappa r) \right] k(r, t) dr$$

A.4

In Chapter 2, the velocity spectrum tensor was defined as

$$\mathcal{E}_{ij}(\underline{\kappa}, t) = \frac{1}{(2\pi)^3} \int_{\mathcal{V}} \mathcal{R}_{ij}(\underline{r}, t) e^{i\underline{\kappa} \cdot \underline{r}} d\underline{r}$$

Using a contraction of indices

$$\mathcal{E}_{ii}(\underline{\kappa}, t) = \frac{1}{(2\pi)^3} \int_{\mathcal{V}} \mathcal{R}_{ii}(\underline{r}, t) e^{i\underline{\kappa} \cdot \underline{r}} d\underline{r}$$

And for isotropic turbulence, as shown in Chapter 4 Part 2

$$\mathcal{R}_{ii}(r, t) = 3f(r) + rf'(r)$$

Proves that \mathcal{R}_{ii} is only a function of r , a scalar quantity.

Following the same steps taken for $T_{ii}(\kappa)$ in Appendix A.2, the following result is obtained,

$$\mathcal{E}_{ii}(\kappa) = \frac{2}{(2\pi)^2} \int_0^\infty r^2 \mathcal{R}_{ii}(r) \frac{\sin(\kappa r)}{\kappa r} dr \quad (11A)$$

It is also possible to relate $\mathcal{E}_{ii}(\kappa)$ to $E(\kappa)$, as shown in alternative derivation for relation between 1D and 3D spectra, obtaining the equation:

$$E(\kappa) = 2\pi\kappa^2 \mathcal{E}_{ii}(\kappa) \quad (12A)$$

Substituting Eq. (11A) into (12A) gives

$$E(\kappa) = \kappa^2 \frac{1}{\pi} \int_0^\infty r^2 \mathcal{R}_{ii}(r) \frac{\sin(\kappa r)}{\kappa r} dr \quad (13A)$$

And substituting Eq. (13A) into (12A) yields

$$E(\kappa) = \frac{\overline{u^2}}{\pi} \int_0^\infty (3f(r) + rf'(r)) \kappa r \sin(\kappa r) dr$$

A.5

$$f(r, t) = e^{-\frac{r^2}{2\lambda_g^2}} \quad (14A)$$

Evaluate 1D spectrum E_{11} using (Chapter 4 Part 5 Eq. (14))

$$E_{11}(\kappa_1, t) = \frac{2}{\pi} \overline{u^2} \int_0^\infty f(r, t) \cos \kappa_1 r \, dr \quad (15A)$$

Substituting Eq. (14A) into (15A) gives

$$E_{11}(\kappa_1, t) = \frac{2}{\pi} \overline{u^2} \int_0^\infty e^{-\frac{r^2}{2\lambda_g^2}} \cos \kappa_1 r \, dr$$

This integral can be reconducted to a differential equation. Differentiate with respect to κ_1

$$\frac{dE_{11}(\kappa_1, t)}{d\kappa_1} = -\frac{2}{\pi} \overline{u^2} \int_0^\infty r e^{-\frac{r^2}{2\lambda_g^2}} \sin \kappa_1 r \, dr$$

Use the substitution:

$$de^{-\frac{r^2}{2\lambda_g^2}} = -\frac{r}{\lambda_g^2} e^{-\frac{r^2}{2\lambda_g^2}} dr$$

To obtain

$$\frac{dE_{11}(\kappa_1, t)}{d\kappa_1} = \frac{2}{\pi} \overline{u^2} \lambda_g^2 \int_0^\infty \sin \kappa_1 r \, de^{-\frac{r^2}{2\lambda_g^2}}$$

Integrate by parts:

$$\frac{dE_{11}(\kappa_1, t)}{d\kappa_1} = \frac{2}{\pi} \overline{u^2} \lambda_g^2 \left\{ \left[\sin(\kappa_1 r) e^{-\frac{r^2}{2\lambda_g^2}} \right]_0^\infty - \kappa_1 \int_0^\infty \cos \kappa_1 r e^{-\frac{r^2}{2\lambda_g^2}} dr \right\}$$

$$\frac{dE_{11}(\kappa_1, t)}{d\kappa_1} = -\frac{2}{\pi} \overline{u^2} \lambda_g^2 \kappa_1 \int_0^\infty \cos \kappa_1 r e^{-\frac{r^2}{2\lambda_g^2}} dr$$

Which is equal to:

$$\frac{dE_{11}(\kappa_1, t)}{d\kappa_1} = -\lambda_g^2 \kappa_1 E_{11}(\kappa_1, t)$$

And using separation of variables yields

$$\frac{dE_{11}(\kappa_1, t)}{E_{11}(\kappa_1, t)} = -\lambda_g^2 \kappa_1 d\kappa_1$$

$$E_{11}(\kappa_1, t) = C \exp\left(-\frac{\lambda_g^2 \kappa_1^2}{2}\right)$$

Where the constant C is found by evaluating

$$E_{11}(0, t) = \frac{2}{\pi} \overline{u^2} \int_0^\infty f(r, t) dr = \frac{2}{\pi} \overline{u^2} \int_0^\infty e^{-\frac{r^2}{2\lambda_g^2}} dr = \frac{\sqrt{2}}{\sqrt{\pi}} \overline{u^2} \lambda_g$$

Such that

$$E_{11}(\kappa_1, t) = \frac{\sqrt{2}}{\sqrt{\pi}} \overline{u^2} \lambda_g \exp\left(-\frac{\lambda_g^2 \kappa_1^2}{2}\right) \quad (16A)$$

Using the relation between 1D and 3D spectra (Chapter 4 Part 5 Appendix A.2)

$$E(\kappa_1) = \frac{\kappa_1^2}{2} \frac{d^2 E_{11}}{d\kappa_1^2} - \frac{\kappa_1}{2} \frac{dE_{11}}{d\kappa_1}$$

And substituting Eq. (16A) gives

$$E(\kappa_1) = \frac{\sqrt{2}}{\sqrt{\pi}} \overline{u^2} \lambda_g \left[\frac{\kappa_1^2}{2} (\lambda_g^4 \kappa_1^2 - \lambda_g^2) \exp\left(-\frac{\lambda_g^2 \kappa_1^2}{2}\right) + \frac{\kappa_1}{2} \kappa_1 \lambda_g^2 \exp\left(-\frac{\lambda_g^2 \kappa_1^2}{2}\right) \right]$$

$$E(\kappa_1) = \frac{\overline{u^2} \lambda_g}{\sqrt{2\pi}} (\lambda_g \kappa_1)^4 \exp\left(-\frac{\lambda_g^2 \kappa_1^2}{2}\right)$$

A.6

$$\begin{aligned} S_{ii}(r, t) &= u_{rms}^3 \frac{1}{r^2} \frac{d}{dr} \left(r^3 \frac{dk}{dr}(r, t) + 4r^2 k(r, t) \right) \\ &= u_{rms}^3 \frac{1}{r^2} \frac{d}{dr} \left(3r^2 \frac{dk}{dr}(r, t) + r^3 \frac{d^2 k}{dr^2}(r, t) + 8rk(r, t) + 4r^2 \frac{dk}{dr}(r, t) \right) \\ S_{ii}(r, t) &= u_{rms}^3 \left(7 \frac{dk}{dr}(r, t) + r \frac{d^2 k}{dr^2}(r, t) + \frac{8}{r} k(r, t) \right) \end{aligned}$$

A.7

$$S_{ii}(r, t) = u_{rms}^3 \left(7 \frac{dk}{dr}(r, t) + r \frac{d^2k}{dr^2}(r, t) + \frac{8}{r} k(r, t) \right) \quad (17A)$$

Taylor expansion for $k(r, t)$ and its derivatives

$$k(r, t) \approx \frac{r^3}{3!} k'''(0, t)$$

$$\frac{dk}{dr}(r, t) \approx 3 \frac{r^2}{3!} k'''(0, t) = \frac{r^2}{2} k'''(0, t)$$

$$\frac{d^2k}{dr^2}(r, t) \approx 2 \frac{r}{2} k'''(0, t) = r k'''(0, t)$$

Substituting these expressions into Eq. (17A) gives

$$S_{ii}(r, t) = u_{rms}^3 \left(7 \frac{r^2}{2} k'''(0, t) + r^2 k'''(0, t) + \frac{4}{3} r^2 k'''(0, t) \right) = \frac{35}{6} r^2 k'''(0, t)$$