

This Hypercard stack was prepared by:
Dennis L. Bricker,
Dept. of Industrial Engineering,
University of Iowa,
Iowa City, Iowa 52242
e-mail: dbricker@icaen.uiowa.edu



Cutting Plane Algorithms for Integer Programming



Cutting-Plane Techniques: From a non-integer optimal solution of the LP relaxation, a constraint is derived and added to the LP, such that the LP solution is eliminated, but NO integer feasible solution is eliminated.



Gomory's Fractional Cut

Dual All-Integer Cut

Gomory's Fractional Cut

Suppose that the optimal LP tableau includes the row

$$\sum_{j=1}^n \alpha_{ij} x_j = \beta_i$$

Suppose that x_k is basic in this row, so that

$$x_k + \sum_{j \notin B} \alpha_{ij} x_j = \beta_i$$

where B = index set of basic variables.

©Dennis Bricker, U. of Iowa, 1997

Notation


$[\alpha_{ij}]$ = integer part of α_{ij}

f_{ij} = fractional part of
 $= \alpha_{ij} - [\alpha_{ij}]$

Examples

$$\left[\frac{5}{4} \right] = 1 \quad \left[\frac{3}{4} \right] = 0$$

$$\left[-\frac{3}{4} \right] = -1$$

 Note that $[a] \leq a$

$[\beta_i]$ = integer part of β_i

f_i = fractional part of
 $= \beta_i - [\beta_i]$

©Dennis Bricker, U. of Iowa, 1997

$$x_k + \sum_{j \notin B} \alpha_{ij} x_j = \beta_i$$

may be written

$$x_k + \sum_{j \notin B} ([\alpha_{ij}] + f_{ij}) x_j = [\beta_i] + f_i$$

$$\Rightarrow x_k - [\beta_i] + \sum_{j \notin B} [\alpha_{ij}] x_j = f_i - \sum_{j \notin B} f_{ij} x_j$$

©Dennis Bricker, U. of Iowa, 1997

A **NECESSARY** condition for x_k & x_j ($j \notin B$) to be integer is that the right-hand-side of

$$x_k - [\beta_i] + \sum_{j \notin B} [\alpha_{ij}] x_j = f_i - \sum_{j \notin B} f_{ij} x_j$$

is integer, i.e.,

$$f_i - \sum_{j \notin B} f_{ij} x_j \in \{\dots -2, -1, 0, 1, 2, 3, \dots\}$$

©Dennis Bricker, U. of Iowa, 1997

However, $f_i < 1$ & $f_{ij}x_j \geq 0$

imply that $f_i - \sum_{j \in B} f_{ij}x_j < 1$

and, indeed, $f_i - \sum_{j \in B} f_{ij}x_j$ must be no greater

than the largest integer < 1 , i.e.,

$$f_i - \sum_{j \in B} f_{ij}x_j \leq 0$$

©Dennis Bricker, U. of Iowa, 1997

Gomory's Fractional Cut

$$f_i - \sum_{j \in B} f_{ij}x_j \leq 0$$

$$\Rightarrow \sum_{j \in B} f_{ij}x_j \geq f_i$$

$$- \sum_{j \in B} f_{ij}x_j \leq -f_i$$

$$- \sum_{j \in B} f_{ij}x_j + S = -f_i$$

*slack
variable*

©Dennis Bricker, U. of Iowa, 1997

Gomory's Fractional Cut

$$-\sum_{j \notin B} f_{ij}x_j + S = -f_i$$

This constraint **MUST** be satisfied by all **INTEGER** feasible solutions of the source row!

However, it is **NOT** satisfied by the current LP solution if $f_i \neq 0!$

(Since $x_j=0$ for $j \notin B$)

©Dennis Bricker, U. of Iowa, 1997

Gomory's Fractional Cut

$$\sum_{j \in B} f_{ij} x_j \geq f_i$$

Example

basic variable $\rightarrow x_4$

row of optimal LP tableau

| | x_1 | x_2 | x_3 | x_4 | x_5 | x_6 | x_7 | RHS | |
|--|---|-------|---------------|-------|-------|---------------|----------------|--------------------|-------------------|
| | 0 | 3 | $\frac{1}{4}$ | 1 | 0 | $\frac{1}{3}$ | $\frac{11}{4}$ | $\frac{21}{5}$ | <i>source row</i> |
| | ↓ | ↓ | ↓ | ↓ | ↓ | ↓ | ↓ | ↓ | |
| | 0 | 0 | $\frac{1}{4}$ | 0 | 0 | $\frac{2}{3}$ | $\frac{3}{4}$ | $\geq \frac{1}{5}$ | |
| | $\frac{1}{4}x_3 + \frac{2}{3}x_6 + \frac{3}{4}x_7 \geq \frac{1}{5}$ | | | | | | | | |

©Dennis Bricker, U. of Iowa, 1997

Example

$$\frac{1}{4}x_3 + \frac{2}{3}x_6 + \frac{3}{4}x_7 \geq \frac{1}{5}$$

If x_3 , x_6 , and x_7 are nonbasic in the current LP optimal tableau, then these variables are ZERO in the basic solution, and the above constraint is violated by the current LP optimal solution!

©Dennis Bricker, U. of Iowa, 1997

Gomory's Cutting-Plane Algorithm

Step 0 Initialization

Solve the LP relaxation of the problem

Step 1 Optimality test

Is the LP solution integer? If so, stop.

Step 2 Cut

Choose a source row (with non-integer right-hand-side) and generate a cut.

Add cut to bottom of tableau

©Dennis Bricker, U. of Iowa, 1997

Step 3**Pivot**

Re-optimize the LP, using the dual simplex algorithm.

Return to step 1.

©Dennis Bricker, U. of Iowa, 1997

All variables (including slack/surplus variables) must be integer.

If original inequality constraint has non-integer coefficients or right-hand-side, multiply both sides by an appropriate positive constant, e.g.

$$\frac{2}{5} x_1 + \frac{4}{3} x_2 \leq \frac{5}{2}$$

multiply both sides by 30

$$\Rightarrow 12 x_1 + 40 x_2 \leq 75$$

©Dennis Bricker, U. of Iowa, 1997

Choice of Source Row

Cuts may be generated using as source row:

- any row in optimal LP tableau which has a non-integer right-hand-side
- a multiple of any row in the LP tableau
- a linear combination of rows from the LP tableau

©Dennis Bricker, U. of Iowa, 1997

Choice of Source Row

While the strength of the cut varies, depending upon one's choice, no rule is known which will guarantee choosing the row yielding the strongest cut.

©Dennis Bricker, U. of Iowa, 1997

Heuristic rules


Choose, as source row, that which has

- 1) $\max_i \{f_i\}$
- 2) $\max_i \left\{ \frac{f_i}{\sum_{j \notin B} f_{ij}} \right\}$
- 3) $\min \left\{ \frac{1}{2} - f_i \right\}$

©Dennis Bricker, U. of Iowa, 1997

$$\begin{aligned} \text{Max } z &= 2x_1 + x_2 \\ \text{s.t. } \quad &x_1 + x_2 \leq 5 \\ &-x_1 + x_2 \leq 0 \\ &6x_1 + 2x_2 \leq 21 \\ &x_1, x_2 \geq 0 \text{ \& integer} \end{aligned}$$

EXAMPLE



©Dennis Bricker, U. of Iowa, 1997

EXAMPLE

Introduce slack variables to convert to equations:

$$\begin{aligned}
 \text{Max } z &= 2x_1 + x_2 \\
 \text{subject to } x_1 + x_2 + x_3 &= 5 \\
 -x_1 + x_2 + x_4 &= 0 \\
 6x_1 + 2x_2 + x_5 &= 21 \\
 x_j &\in \{0, 1, 2, 3, \dots\}
 \end{aligned}$$

©Dennis Bricker, U. of Iowa, 1997

EXAMPLE

| | -Z | x ₁ | x ₂ | x ₃ | x ₄ | x ₅ | rhs |
|-----------------------|----|----------------|----------------|----------------|----------------|----------------|-----------------|
| optimal LP tableau | 1 | 0 | 0 | $-\frac{1}{2}$ | 0 | $-\frac{1}{4}$ | $-\frac{31}{4}$ |
| | 0 | 1 | 0 | $-\frac{1}{2}$ | 0 | $\frac{1}{4}$ | $\frac{11}{4}$ |
| | 0 | 0 | 1 | $\frac{3}{2}$ | 0 | $-\frac{1}{4}$ | $\frac{9}{4}$ |
| | 0 | 0 | 0 | -2 | 1 | $\frac{1}{2}$ | $\frac{1}{2}$ |

©Dennis Bricker, U. of Iowa, 1997

ANY of these rows could serve as the
SOURCE row for a cut:

| | source row | ⇒ | cut |
|-------|---|---|--|
| x_1 | $-\frac{1}{2}x_3 + \frac{1}{4}x_5 = \frac{11}{4}$ | | $\frac{1}{2}x_3 + \frac{1}{4}x_5 \geq \frac{3}{4}$ |
| x_2 | $+\frac{3}{2}x_3 - \frac{1}{4}x_5 = \frac{0}{4}$ | | $\frac{1}{2}x_3 + \frac{3}{4}x_5 \geq \frac{1}{4}$ |
| | $-2x_3 + x_4 + \frac{1}{2}x_5 = \frac{1}{2}$ | | $\frac{1}{2}x_5 \geq \frac{1}{2}$ |

©Dennis Bricker, U. of Iowa, 1997

Graphical Representation of Cuts in X_1X_2 -plane

substitute


$$\begin{cases} x_3 = 5 - x_1 - x_2 \\ x_5 = 21 - 6x_1 - 2x_2 \end{cases}$$

| | ⇒ | cut |
|--|---|-----------------------|
| $\frac{1}{2}x_3 + \frac{1}{4}x_5 \geq \frac{3}{4}$ | | $2x_1 + x_2 \leq 7$ |
| $\frac{1}{2}x_3 + \frac{3}{4}x_5 \geq \frac{1}{4}$ | | $5x_1 + 2x_2 \leq 18$ |
| $\frac{1}{2}x_5 \geq \frac{1}{2}$ | | $6x_1 + 3x_2 \leq 20$ |




*click mouse on
cut to see effect*


©Dennis Bricker, U. of Iowa, 1997



©Dennis Bricker, U. of Iowa, 1997



©Dennis Bricker, U. of Iowa, 1997



©Dennis Bricker, U. of Iowa, 1997

Dropping Cuts from Tableau

Each cut adds a new row & a new column (slack variable) to the tableau...

If ALL cuts are kept until the algorithm terminates, the tableau becomes so large as to be "unwieldy"!

When a cut is no longer "useful", it would be advantageous to be able to delete that cut.



©Dennis Bricker, U. of Iowa, 1997

Dropping Cuts from Tableau


When a cut is added to the tableau, & the dual simplex pivot removes its slack variable from the basis, the cut is a "tight" constraint, i.e., its slack variable is zero.

If a cut's slack variable re-enters the basis at a later iteration, then the cut has become inactive and may then be dropped from the tableau.

©Dennis Bricker, U. of Iowa, 1997

EXAMPLE

$$\begin{aligned}
 &\text{Max } z = 2x_1 + x_2 \\
 &\text{s.t.} \quad x_1 + x_2 \leq 5 \\
 &\quad \quad -x_1 + x_2 \leq 0 \\
 &\quad \quad 6x_1 + 2x_2 \leq 21 \\
 &\quad \quad x_1, x_2 \geq 0 \text{ \& integer}
 \end{aligned}$$



Initial Optimal LP tableau

| |
|--------------------|
| Current LP Tableau |
|--------------------|

| z | 1 | 2 | 3 | 4 | 5 | B |
|---|---|---|------|---|-------|-------|
| 1 | 0 | 0 | -0.5 | 0 | -0.25 | -7.75 |
| 0 | 0 | 1 | 1.5 | 0 | -0.25 | 2.25 |
| 0 | 0 | 0 | -2 | 1 | 0.5 | 0.5 |
| 0 | 1 | 0 | -0.5 | 0 | 0.25 | 2.75 |

Variables:

(Negative of) objective function value: z
 Original structural variables: 1 2
 Original slack/surplus variables: 3 4 5
 Slack variables for cuts:

©Dennis Bricker, U. of Iowa, 1997

The rows having non-integer right-hand-side are 2 3 4

Source row is # 2

| i | 2 | 3 | 5 | 6 | rhs |
|------------|---|------|-------|---|-------|
| Source row | 1 | 1.5 | -0.25 | 0 | 2.25 |
| Cut | 0 | -0.5 | -0.75 | 1 | -0.25 |

(X[6] (= slack variable for new cut) is basic but < 0)

The cut which is added is (in terms of original variables):

| 1 | 2 | b |
|---|---|------|
| 5 | 2 | ≤ 18 |

©Dennis Bricker, U. of Iowa, 1997

| |
|--------------------|
| Current LP Tableau |
|--------------------|

| z | 1 | 2 | 3 | 4 | 5 | 6 | B |
|---|---|---|------|---|-------|---|-------|
| 1 | 0 | 0 | -0.5 | 0 | -0.25 | 0 | -7.75 |
| 0 | 0 | 1 | 1.5 | 0 | -0.25 | 0 | 2.25 |
| 0 | 0 | 0 | -2 | 1 | 0.5 | 0 | 0.5 |
| 0 | 1 | 0 | -0.5 | 0 | 0.25 | 0 | 2.75 |
| 0 | 0 | 0 | -0.5 | 0 | -0.75 | 1 | -0.25 |

← cut

Variables:

(Negative of) objective function value: z
 Original structural variables: 1 2
 Original slack/surplus variables: 3 4 5
 Slack variables for cuts: 6

**Tableau is now
 primal infeasible
 (but dual feasible!)**

©Dennis Bricker, U. of Iowa, 1997

| |
|--------------------|
| Solving current LP |
|--------------------|

Performing dual simplex pivot in row 5

Potential pivot columns: X{3 5}

| i | 3 | 5 |
|-------------|------|-------|
| Rel. Profit | -0.5 | -0.25 |
| Subs. rate | -0.5 | -0.75 |
| Ratio | 1 | 0.333 |

Minimum ratio is in column 5,
 which is selected as pivot column

©Dennis Bricker, U. of Iowa, 1997

Current LP Tableau

| z | 1 | 2 | 3 | 4 | 5 | 6 | B |
|---|---|---|------|---|-------|---|-------|
| 1 | 0 | 0 | -0.5 | 0 | -0.25 | 0 | -7.75 |
| 0 | 0 | 1 | 1.5 | 0 | -0.25 | 0 | 2.25 |
| 0 | 0 | 0 | -2 | 1 | 0.5 | 0 | 0.5 |
| 0 | 1 | 0 | -0.5 | 0 | -0.25 | 0 | 2.75 |
| 0 | 0 | 0 | -0.5 | 0 | -0.75 | 1 | -0.25 |

↑
pivot
column

©Dennis Bricker, U. of Iowa, 1997

Current LP Tableau

| z | 1 | 2 | 3 | 4 | 5 | 6 | B |
|---|---|---|--------|---|---|--------|-------|
| 1 | 0 | 0 | -0.333 | 0 | 0 | -0.333 | -7.67 |
| 0 | 0 | 1 | 1.67 | 0 | 0 | -0.333 | 2.33 |
| 0 | 0 | 0 | -2.33 | 1 | 0 | 0.667 | 0.333 |
| 0 | 1 | 0 | -0.667 | 0 | 0 | 0.333 | 2.67 |
| 0 | 0 | 0 | 0.667 | 0 | 1 | -1.33 | 0.333 |

Variables:

(Negative of) objective function value: z
 Original structural variables: 1 2
 Original slack/surplus variables: 3 4 5
 Slack variables for cuts: 6

©Dennis Bricker, U. of Iowa, 1997

The rows having non-integer right-hand-side are 2 3 4 5

Source row is # 2

| i | 2 | 3 | 6 | 7 | rhs |
|------------|---|--------|--------|---|--------|
| Source row | 1 | 1.67 | -0.333 | 0 | 2.33 |
| Cut: | 0 | -0.667 | -0.667 | 1 | -0.333 |

(X[7] (= slack variable for new cut) is basic but < 0)

The cut which is added is (in terms of original variables):

| | | |
|---|---|-----------|
| 1 | 2 | b |
| 4 | 2 | ≤ 15 |

©Dennis Bricker, U. of Iowa, 1997

Current LP Tableau

| z | 1 | 2 | 3 | 4 | 5 | 6 | 7 | B |
|---|---|---|--------|---|---|--------|---|--------|
| 1 | 0 | 0 | -0.333 | 0 | 0 | -0.333 | 0 | -7.67 |
| 0 | 0 | 1 | 1.67 | 0 | 0 | -0.333 | 0 | 2.33 |
| 0 | 0 | 0 | -2.33 | 1 | 0 | 0.667 | 0 | 0.333 |
| 0 | 1 | 0 | -0.667 | 0 | 0 | 0.333 | 0 | 2.67 |
| 0 | 0 | 0 | 0.667 | 0 | 1 | -1.33 | 0 | 0.333 |
| 0 | 0 | 0 | -0.667 | 0 | 0 | -0.667 | 1 | -0.333 |

← cut

Variables:

(Negative of) objective function value: z
 Original structural variables: 1 2
 Original slack/surplus variables: 3 4 5
 Slack variables for cuts: 6 7

©Dennis Bricker, U. of Iowa, 1997

| |
|--------------------|
| Solving current LP |
|--------------------|

Performing dual simplex pivot in row 6

Potential pivot columns: X[3 6]

| i | 3 | 6 |
|-------------|--------|--------|
| Rel. Profit | -0.333 | -0.333 |
| Subs. rate | -0.667 | -0.667 |
| Ratio | 0.5 | 0.5 |

Minimum ratio is in column 3,
which is selected as pivot column

©Dennis Bricker, U. of Iowa, 1997

| |
|--------------------|
| Current LP Tableau |
|--------------------|

| z | 1 | 2 | 3 | 4 | 5 | 6 | 7 | B |
|---|---|---|--------|---|---|--------|---|--------|
| 1 | 0 | 0 | -0.333 | 0 | 0 | -0.333 | 0 | -7.67 |
| 0 | 0 | 1 | 1.67 | 0 | 0 | -0.333 | 0 | 2.33 |
| 0 | 0 | 0 | -2.33 | 1 | 0 | 0.667 | 0 | 0.333 |
| 0 | 1 | 0 | -0.667 | 0 | 0 | 0.333 | 0 | 2.67 |
| 0 | 0 | 0 | -0.667 | 0 | 1 | -1.33 | 0 | 0.333 |
| 0 | 0 | 0 | -0.667 | 0 | 0 | -0.667 | 1 | -0.333 |

↑
pivot
column

©Dennis Bricker, U. of Iowa, 1997

| |
|--------------------|
| Current LP Tableau |
|--------------------|

| z | 1 | 2 | 3 | 4 | 5 | 6 | 7 | B |
|---|---|---|---|---|---|----|------|------|
| 1 | 0 | 0 | 0 | 0 | 0 | 0 | -0.5 | -7.5 |
| 0 | 0 | 1 | 0 | 0 | 0 | -2 | 2.5 | 1.5 |
| 0 | 0 | 0 | 0 | 1 | 0 | 3 | -3.5 | 1.5 |
| 0 | 1 | 0 | 0 | 0 | 0 | 1 | -1 | 3 |
| 0 | 0 | 0 | 0 | 0 | 1 | -2 | 1 | 0 |
| 0 | 0 | 0 | 1 | 0 | 0 | 1 | -1.5 | 0.5 |

Variables:

(Negative of) objective function value: z
 Original structural variables: 1 2
 Original slack/surplus variables: 3 4 5
 Slack variables for cuts: 6 7

©Dennis Bricker, U. of Iowa, 1997

The rows having non-integer right-hand-side are 2 3 6

Source row is # 2

| i | 2 | 6 | 7 | 8 | rhs |
|------------|---|----|------|---|------|
| Source row | 1 | -2 | 2.5 | 0 | 1.5 |
| Cut: | 0 | 0 | -0.5 | 1 | -0.5 |

(X[8] (= slack variable for new cut) is basic but < 0)

The cut which is added is (in terms of original variables):

| | | |
|---|---|----------|
| 1 | 2 | b |
| 2 | 1 | ≤ 7 |

©Dennis Bricker, U. of Iowa, 1997

| |
|--------------------|
| Current LP Tableau |
|--------------------|

| z | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | B |
|---|---|---|---|---|---|----|------|---|------|
| 1 | 0 | 0 | 0 | 0 | 0 | 0 | -0.5 | 0 | -7.5 |
| 0 | 0 | 1 | 0 | 0 | 0 | -2 | 2.5 | 0 | 1.5 |
| 0 | 0 | 0 | 0 | 1 | 0 | 3 | -3.5 | 0 | 1.5 |
| 0 | 1 | 0 | 0 | 0 | 0 | 1 | -1 | 0 | 3 |
| 0 | 0 | 0 | 0 | 0 | 1 | -2 | 1 | 0 | 0 |
| 0 | 0 | 0 | 1 | 0 | 0 | 1 | -1.5 | 0 | 0.5 |
| 0 | 0 | 0 | 0 | 0 | 0 | 0 | -0.5 | 1 | -0.5 |

← cut

Variables:

(Negative of) objective function value: z
 Original structural variables: 1 2
 Original slack/surplus variables: 3 4 5
 Slack variables for cuts: 6 7 8

©Dennis Bricker, U. of Iowa, 1997

| |
|--------------------|
| Solving current LP |
|--------------------|

Performing dual simplex pivot in row 7

Potential pivot columns: X[7]

| | |
|-------------|------|
| i: | 7 |
| Rel. Profit | -0.5 |
| Subs. rate | -0.5 |
| Ratio | 1 |

Minimum ratio is in column 7,
 which is selected as pivot column

Resulting solution is again infeasible (variable < 0)

©Dennis Bricker, U. of Iowa, 1997

Current LP Tableau

| z | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | B |
|---|---|---|---|---|---|----|------|---|------|
| 1 | 0 | 0 | 0 | 0 | 0 | 0 | -0.5 | 0 | -7.5 |
| 0 | 0 | 1 | 0 | 0 | 0 | -2 | 2.5 | 0 | 1.5 |
| 0 | 0 | 0 | 0 | 1 | 0 | 3 | -3.5 | 0 | 1.5 |
| 0 | 1 | 0 | 0 | 0 | 0 | 1 | -1 | 0 | 3 |
| 0 | 0 | 0 | 0 | 0 | 1 | -2 | 1 | 0 | 0 |
| 0 | 0 | 0 | 1 | 0 | 0 | 1 | -1.5 | 0 | 0.5 |
| 0 | 0 | 0 | 0 | 0 | 0 | 0 | -0.5 | 1 | -0.5 |

©Dennis Bricker, U. of Iowa, 1997

| z | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | B |
|---|---|---|---|---|---|----|---|----|----|
| 1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | -1 | -7 |
| 0 | 0 | 1 | 0 | 0 | 0 | -2 | 0 | 5 | -1 |
| 0 | 0 | 0 | 0 | 1 | 0 | 3 | 0 | -7 | 5 |
| 0 | 1 | 0 | 0 | 0 | 0 | 1 | 0 | -2 | 4 |
| 0 | 0 | 0 | 0 | 0 | 1 | -2 | 0 | 2 | -1 |
| 0 | 0 | 0 | 1 | 0 | 0 | 1 | 0 | -3 | 2 |
| 0 | 0 | 0 | 0 | 0 | 0 | 0 | 1 | -2 | 1 |

As a result of the previous dual simplex pivot, the right-hand-side of the new row becomes positive, but further dual simplex pivots are necessary, because negative numbers have appeared in other rows!

©Dennis Bricker, U. of Iowa, 1997

| z | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | B |
|---|---|---|---|---|---|----|---|----|----|
| 1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | -1 | -7 |
| 0 | 0 | 1 | 0 | 0 | 0 | -2 | 0 | 5 | -1 |
| 0 | 0 | 0 | 0 | 1 | 0 | 3 | 0 | -7 | 5 |
| 0 | 1 | 0 | 0 | 0 | 0 | 1 | 0 | -2 | 4 |
| 0 | 0 | 0 | 0 | 0 | 1 | -2 | 0 | 2 | -1 |
| 0 | 0 | 0 | 1 | 0 | 0 | 1 | 0 | -3 | 2 |
| 0 | 0 | 0 | 0 | 0 | 0 | 0 | 1 | -2 | 1 |

Next pivot row should be either row 2 or row 5.

Performing dual simplex pivot in row 2

Potential pivot columns: X[6]

| | |
|-------------|----|
| i | 6 |
| Rel. Profit | 0 |
| Subs. rate | -2 |
| Ratio | 0 |

Minimum ratio is in column 6,
which is selected as pivot column

©Dennis Bricker, U. of Iowa, 1997

| z | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | B |
|---|---|---|---|---|---|----|---|----|----|
| 1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | -1 | -7 |
| 0 | 0 | 1 | 0 | 0 | 0 | -2 | 0 | 5 | -1 |
| 0 | 0 | 0 | 0 | 1 | 0 | 3 | 0 | -7 | 5 |
| 0 | 1 | 0 | 0 | 0 | 0 | 1 | 0 | -2 | 4 |
| 0 | 0 | 0 | 0 | 0 | 1 | -2 | 0 | 2 | -1 |
| 0 | 0 | 0 | 1 | 0 | 0 | 1 | 0 | -3 | 2 |
| 0 | 0 | 0 | 0 | 0 | 0 | 0 | 1 | -2 | 1 |

©Dennis Bricker, U. of Iowa, 1997

| |
|--------------------|
| Current LP Tableau |
|--------------------|

| z | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | B |
|---|---|------|---|---|---|---|---|------|-----|
| 1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | -1 | -7 |
| 0 | 0 | -0.5 | 0 | 0 | 0 | 1 | 0 | -2.5 | 0.5 |
| 0 | 0 | 1.5 | 0 | 1 | 0 | 0 | 0 | 0.5 | 3.5 |
| 0 | 1 | 0.5 | 0 | 0 | 0 | 0 | 0 | 0.5 | 3.5 |
| 0 | 0 | -1 | 0 | 0 | 1 | 0 | 0 | -3 | 0 |
| 0 | 0 | 0.5 | 1 | 0 | 0 | 0 | 0 | -0.5 | 1.5 |
| 0 | 0 | 0 | 0 | 0 | 0 | 0 | 1 | -2 | 1 |

Variables:

(Negative of) objective function value: z
 Original structural variables: 1 2
 Original slack/surplus variables: 3 4 5
 Slack variables for cuts: 6 7 8

©Dennis Bricker, U. of Iowa, 1997

The rows having non-integer right-hand-side are 2 3 4 6

From which row do you wish to generate the cut?

□:

2

The cut which is added is (in terms of original variables):

| | | |
|---|---|-----|
| 1 | 2 | b |
| 1 | 0 | ≤ 3 |

Source row is # 2

| | | | | | | |
|-------------|-------|------|---|------|---|------|
| i: | _____ | 2 | 6 | 8 | 9 | rhs |
| Source row: | | -0.5 | 1 | -2.5 | 0 | 0.5 |
| Cut: | | -0.5 | 0 | -0.5 | 1 | -0.5 |

(X[9] (= slack variable for new cut) is basic but < 0)

©Dennis Bricker, U. of Iowa, 1997

| |
|--------------------|
| Current LP Tableau |
|--------------------|

| z | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | B |
|---|---|------|---|---|---|---|---|------|---|------|
| 1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | -1 | 0 | -7 |
| 0 | 0 | -0.5 | 0 | 0 | 0 | 1 | 0 | -2.5 | 0 | 0.5 |
| 0 | 0 | 1.5 | 0 | 1 | 0 | 0 | 0 | 0.5 | 0 | 3.5 |
| 0 | 1 | 0.5 | 0 | 0 | 0 | 0 | 0 | 0.5 | 0 | 3.5 |
| 0 | 0 | -1 | 0 | 0 | 1 | 0 | 0 | -3 | 0 | 0 |
| 0 | 0 | 0.5 | 1 | 0 | 0 | 0 | 0 | -0.5 | 0 | 1.5 |
| 0 | 0 | 0 | 0 | 0 | 0 | 0 | 1 | -2 | 0 | 1 |
| 0 | 0 | -0.5 | 0 | 0 | 0 | 0 | 0 | -0.5 | 1 | -0.5 |

Variables:

(Negative of) objective function value: z
 Original structural variables: 1 2
 Original slack/surplus variables: 3 4 5
 Slack variables for cuts: 6 7 8 9

©Dennis Bricker, U. of Iowa, 1997

| |
|--------------------|
| Solving current LP |
|--------------------|

Performing dual simplex pivot in row 8

Potential pivot columns: X[2 8]

| | | |
|-------------|------|------|
| i: | 2 | 8 |
| Rel. Profit | 0 | -1 |
| Subs. rate | -0.5 | -0.5 |
| Ratio | 0 | 2 |

Minimum ratio is in column 2,
 which is selected as pivot column

©Dennis Bricker, U. of Iowa, 1997

| z | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | B |
|---|---|------|---|---|---|---|---|------|---|------|
| 1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | -1 | 0 | -7 |
| 0 | 0 | -0.5 | 0 | 0 | 0 | 1 | 0 | -2.5 | 0 | 0.5 |
| 0 | 0 | 1.5 | 0 | 1 | 0 | 0 | 0 | 0.5 | 0 | 3.5 |
| 0 | 1 | 0.5 | 0 | 0 | 0 | 0 | 0 | 0.5 | 0 | 3.5 |
| 0 | 0 | -1 | 0 | 0 | 1 | 0 | 0 | -3 | 0 | 0 |
| 0 | 0 | 0.5 | 1 | 0 | 0 | 0 | 0 | -0.5 | 0 | 1.5 |
| 0 | 0 | 0 | 0 | 0 | 0 | 0 | 1 | -2 | 0 | 1 |
| 0 | 0 | -0.5 | 0 | 0 | 0 | 0 | 0 | -0.5 | 1 | -0.5 |

©Dennis Bricker, U. of Iowa, 1997

Current LP Tableau

| z | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | 9 | B |
|---|---|---|---|---|---|---|---|----|----|----|
| 1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | -1 | 0 | -7 |
| 0 | 0 | 0 | 0 | 0 | 0 | 1 | 0 | -2 | -1 | 1 |
| 0 | 0 | 0 | 0 | 1 | 0 | 0 | 0 | -1 | 3 | 2 |
| 0 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 3 |
| 0 | 0 | 0 | 0 | 0 | 1 | 0 | 0 | -2 | -2 | 1 |
| 0 | 0 | 0 | 1 | 0 | 0 | 0 | 0 | -1 | 1 | 1 |
| 0 | 0 | 0 | 0 | 0 | 0 | 0 | 1 | -2 | 0 | 1 |
| 0 | 0 | 1 | 0 | 0 | 0 | 0 | 0 | 1 | -2 | 1 |

All variables are integer!

Variables:

- (Negative of) objective function value: z
- Original structural variables: 1 2
- Original slack/surplus variables: 3 4 5
- Slack variables for cuts: 6 7 8 9

©Dennis Bricker, U. of Iowa, 1997

Current List of Cuts

| # | 1 | 2 | b |
|----|---|---|------|
| 1) | 5 | 2 | ≤ 18 |
| 2) | 4 | 2 | ≤ 15 |
| 3) | 2 | 1 | ≤ 7 |
| 4) | 1 | 0 | ≤ 3 |

