


Pre-Calculus Facility Location Problems

This Hypercard stack was prepared by:
 Dennis L. Bricker,
 Dept. of Industrial Engineering,
 University of Iowa,
 Iowa City, Iowa 52242
 e-mail: dbricker@icaen.uiowa.edu

- "The Stream-side Tavern" Location Problem
- Locating a Market for Three Villages
- Locating a Market for more than 3 Villages (Weber's Problem)
- The Ice-Cream Vendor Problem
- The Firehouse Problem
- The LifeGuard Problem

©Dennis Bricker, U. of Iowa, 1997

The Greek Streamside Tavern




Villagers returning home after working in the communal field will walk to the stream to fill a bucket with water, and then walk to the village.

Where would be the best location for a tavern along the stream? (Assume X is the location which minimizes the total distance walked!)




©Dennis Bricker, U. of Iowa, 1997

Minimize $f(X) = d(F,X) + d(X,V)$




©Dennis Bricker, U. of Iowa, 1997

Geometric solution:




©Dennis Bricker, U. of Iowa, 1997



©Dennis Bricker, U. of Iowa, 1997

Locating a marketplace for 3 villages




i.e., select X to minimize $d(X,A) + d(X,B) + d(X,C)$




©Dennis Bricker, U. of Iowa, 1997

Geometric solution




©Dennis Bricker, U. of Iowa, 1997

Find point X' so that $\triangle BX'X$ is equilateral.



©Dennis Bricker, U. of Iowa, 1997

$\triangle BXA \cong \triangle BX'C'$
 since $\angle C'BX' = \angle ABX$
 $C'B = AB$
 & $BX' = BX$



$$f(X) = d(A,X) + d(B,X) + d(C,X)$$


$$= d(C',X') + d(X',X) + d(X,C)$$

$$= \text{length of the path } C'X'XC$$

$\therefore f(X)$ is minimized if $C'X'XC$ are colinear,
 i.e., if X is on the line segment $C'C$


©Dennis Bricker, U. of Iowa, 1997

X is therefore found by constructing C' so that $\triangle ABC'$ is equilateral.



©Dennis Bricker, U. of Iowa, 1997

Next, point A' is constructed so that $\triangle A'BC$ is equilateral.




By symmetry of argument, X must be on the line segment $A'A$.

Therefore, X must lie at the intersection of $C'C$ and $A'A$

©Dennis Bricker, U. of Iowa, 1997

Locating Marketplace for $n > 3$ Villages:



The geometric construction for the case $n=3$ does not generalize for $n > 3$


(*"Weber's Problem"*)



©Dennis Bricker, U. of Iowa, 1997


Suppose that the "weight" at village i is w_i

We wish to minimize $f(X) = \sum_{i=1}^n w_i d(A_i, X)$



©Dennis Bricker, U. of Iowa, 1997

An "analogue computer":




Varignon Frame

Cut holes in a table top on which a map is drawn.

©Dennis Bricker, U. of Iowa, 1997


Tie together in a knot ("X") n strings of equal length L



Attach a bucket with weight w_i at the end of string i .


©Dennis Bricker, U. of Iowa, 1997

The system of strings & weights is released and comes to equilibrium.



©Dennis Bricker, U. of Iowa, 1997

Potential Energy of System is $P.E. = \sum_{i=1}^n W_i h_i$



©Dennis Bricker, U. of Iowa, 1997

$$P.E. = \sum_{i=1}^n W_i h_i = \sum_{i=1}^n (H-L) W_i + W_i d(A_i, X)$$

$$= \underbrace{\sum_{i=1}^n (H-L) W_i}_{constant} + \underbrace{\sum_{i=1}^n W_i d(A_i, X)}_{f(X)}$$

The X which minimizes the potential energy will minimize our objective function!



©Dennis Bricker, U. of Iowa, 1997




An ice cream vendor wants to locate his cart along a beach, minimizing the total distance from the persons on the beach to his cart.



©Dennis Bricker, U. of Iowa, 1997


Consider the beach to be a line, with points representing groups of people: $A_i =$ location of group #i
 $W_i =$ # of persons in group #i



His objective: Minimize $f(x) = \sum_i w_i |x - A_i|$

What is the nature of this objective function, $f(x)$?

©Dennis Bricker, U. of Iowa, 1997



If x lies in the interval $[A_i, A_{i+1}]$,

$$f(x) = \sum_{j \leq i} w_j (x - A_j) + \sum_{j > i} w_j (A_j - x)$$

$$= \sum_{j \leq i} w_j x - \sum_{j \leq i} w_j A_j + \sum_{j > i} w_j A_j - \sum_{j > i} w_j x$$

$$= x \left(\sum_{j \leq i} w_j - \sum_{j > i} w_j \right) + \underbrace{\left(\sum_{j > i} w_j A_j - \sum_{j \leq i} w_j A_j \right)}_{\text{a constant!}}$$

©Dennis Bricker, U. of Iowa, 1997


We have seen that on the interval

$$f(x) = x \left(\sum_{j \leq i} w_j - \sum_{j > i} w_j \right) + (\text{constant})$$

This is a linear function, with slope $\sum_{j \leq i} w_j - \sum_{j > i} w_j$

Therefore, the function is increasing if $\sum_{j \leq i} w_j > \sum_{j > i} w_j$


and decreasing if $\sum_{j \leq i} w_j < \sum_{j > i} w_j$



©Dennis Bricker, U. of Iowa, 1997

That is, the function is increasing if $\sum_{j \leq i} w_j > \sum_{j > i} w_j$

(i.e., there are more people to the left.)



and decreasing if $\sum_{j \leq i} w_j < \sum_{j > i} w_j$


(i.e., there are more people to the right.)

So the ice cream vendor should walk along the beach until half of the bathers are behind him, and half in front of him.

©Dennis Bricker, U. of Iowa, 1997

Note that this means that the optimal solution is at the **median** of the population (not the **mean**, as one might have suspected!)

(For this reason, mini-sum location problems are referred to as **median** problems.)


 Question: what happens to the optimal location if, during the afternoon, a group of bathers to the right of the cart decide to move 100 meters further to the right?



©Dennis Bricker, U. of Iowa, 1997

The problem then is equivalent to finding the smallest circle which circumscribes the three points A, B, and C.

The optimal location is then at the **center** of the circle.




This easily generalizes to n villages, where n > 3.

*For this reason, such "minimax" location problems are referred to as **center** problems!*

©Dennis Bricker, U. of Iowa, 1997


Again we represent the beach by a line, with points indicating the position of the swimmers:



If his objective is to minimize $f(x)$, where

$$f(x) = \text{maximum}_{1 \leq i \leq n} |x - A_i|$$

then the optimal location is the **center** of the smallest circle covering the n points.

 Question: what happens to his optimal location if group #i decides to move to the right? What if $i=n$?

©Dennis Bricker, U. of Iowa, 1997

By definition of absolute value:

$$|x - A_i| = \text{maximum} \{ (x - A_i), (A_i - x) \}$$

So our problem is


$$\text{minimize}_x [\text{maximum}_{1 \leq i \leq n} \{ (x - A_i), (A_i - x) \}]$$

©Dennis Bricker, U. of Iowa, 1997

The Firehouse Location Problem

Suppose that a firehouse is to be located so as to serve three villages. The "best" location is considered to be that which would minimize the longest possible delay in reaching the scene of a fire:


Minimize $g(x)$ where $g(x) = \text{maximum} \{ d(x,A), d(x,B), d(x,C) \}$



$g(x) \leq z \iff A, B, \& C$ are contained within a circle of radius z with center x

©Dennis Bricker, U. of Iowa, 1997

The Life-Guard Problem




The lifeguard wants to locate his chair in a location which will minimize, not the sum of distances to the swimmers, but the distance to the farthest swimmer.



©Dennis Bricker, U. of Iowa, 1997

Suppose that he wants to weight the distances, according to the number of persons in the group:



Let his objective function (to be minimized) be defined by

$$f(x) = \text{maximum}_{1 \leq i \leq n} w_i |x - A_i|$$

in APL: $\Gamma / W \times | X - A$

©Dennis Bricker, U. of Iowa, 1997

$$\text{minimize}_x \{ \text{maximum}_{1 \leq i \leq n} w_i |x - A_i| \}$$

We can transform this optimization problem to an LP by introducing a new variable, z :

$$\begin{aligned} &\text{minimize}_{x,z} z && \text{subject to } z \geq w_i |x - A_i| && \text{for } i=1, \dots, n \\ \Rightarrow &\text{minimize}_{x,z} z && \text{subject to } \begin{cases} z \geq w_i (x - A_i) & \text{for } i=1, \dots, n \\ z \geq w_i (A_i - x) & \text{for } i=1, \dots, n \end{cases} \\ \Rightarrow &\text{minimize}_{x,z} z && \text{subject to } \begin{cases} x \leq A_i + \frac{z}{w_i} & \text{for } i=1, \dots, n \\ x \geq A_i - \frac{z}{w_i} & \text{for } i=1, \dots, n \end{cases} \end{aligned}$$

©Dennis Bricker, U. of Iowa, 1997

The condition for a feasible solution to the constraints

$$\begin{cases} x \leq A_i + \frac{z}{w_i} & \text{for } i=1, \dots, n \\ x \geq A_i - \frac{z}{w_i} & \text{for } i=1, \dots, n \end{cases}$$

is, of course, that every upper bound of x be greater than (or equal to) every lower bound, that is,

$$A_i - \frac{z}{w_i} \leq A_j + \frac{z}{w_j} \quad \text{for every pair } i \& j$$

or, solving these inequalities for z,

$$A_i - A_j \leq \frac{z}{w_j} + \frac{z}{w_i} \Rightarrow \frac{A_i - A_j}{\frac{1}{w_j} + \frac{1}{w_i}} \leq z \quad \text{for all } i \& j$$

©Dennis Bricker, U. of Iowa, 1997

Therefore, we wish to choose the smallest z for which the

$$\text{constraints } \frac{A_i - A_j}{\frac{1}{w_j} + \frac{1}{w_i}} \leq z \quad \text{are feasible}$$

Clearly, then, the optimal z will be the greatest of its lower bounds, i.e.,

$$z = \text{maximum}_{i,j} \left[\frac{A_i - A_j}{\frac{1}{w_j} + \frac{1}{w_i}} \right]$$

What is then the optimal value for the location, x?

©Dennis Bricker, U. of Iowa, 1997

Suppose that the greatest lower bound for z occurs for the pair of points i^* and j^* . Then

$$z = \frac{A_i - A_j}{\frac{1}{w_j} + \frac{1}{w_i}} \Rightarrow A_j - \frac{z}{w_j} \leq x \leq A_i + \frac{z}{w_i}$$

is the "tight" constraint,

$$\text{that is, } A_j - \frac{z}{w_j} = x = A_i + \frac{z}{w_i}$$

and, substituting the optimal value for z, we get

$$x = A_j - \frac{1}{w_j} \left[\frac{A_i - A_j}{\frac{1}{w_j} + \frac{1}{w_i}} \right] = \frac{A_i w_i + A_j w_j}{w_i + w_j}$$

©Dennis Bricker, U. of Iowa, 1997

Notice that if the weights are equal, e.g., $w_i = 1$ for all j, then the optimal value is

$$z = \text{maximum}_{i,j} \left[\frac{A_i - A_j}{\frac{1}{w_j} + \frac{1}{w_i}} \right] = \text{maximum}_{i,j} \left[\frac{A_i - A_j}{2} \right] = \frac{A_n - A_1}{2}$$

(i.e., $i^*=n$ and $j^*=1$)

and the optimal location is

$$x = \frac{A_i w_i + A_j w_j}{w_i + w_j} = \frac{A_n w_n + A_1 w_1}{w_n + w_1} = \frac{A_n + A_1}{2}$$

(i.e., at the midpoint of the interval $[A_1, A_n]$, the *center*) which is what we would expect.

©Dennis Bricker, U. of Iowa, 1997

What is an APL expression for the optimal value in the weighted case, i.e.,

$$z = \text{maximum}_{i,j} \left[\frac{A_i - A_j}{\frac{1}{w_j} + \frac{1}{w_i}} \right] \quad ?$$

$$z = \text{maximum}_{i,j} \left[\frac{A_i - A_j}{\frac{1}{w_j} + \frac{1}{w_i}} \right]$$

would be written in APL as:

$$z \leftarrow \uparrow / , (A \circ . - A) \div (\div w \circ . + w)$$

How could we write the expression for the optimal pair of indices, i^* and j^* , used to maximize above?

©Dennis Bricker, U. of Iowa, 1997

©Dennis Bricker, U. of Iowa, 1997

The vector consisting of i^* and j^* can be found by first locating the maximum value in the "raveled" matrix, i.e.,

$$ij \leftarrow \nu \uparrow \uparrow / \nu \leftarrow , (A \circ . - A) \div (\div w \circ . + w)$$

and then using the "base" function to convert to row number, column number:

$$ij \leftarrow 1 + (2\rho\rho A) \perp^{-1} + ij$$

which could be combined into a single expression:

$$ij \leftarrow 1 + (2\rho\rho A) \perp^{-1} + \nu \uparrow \uparrow / \nu \leftarrow , (A \circ . - A) \div (\div w \circ . + w)$$

©Dennis Bricker, U. of Iowa, 1997

Then the optimal location, i.e.,

$$x = \frac{A_i w_i + A_j w_j}{w_i + w_j}$$

would be written in APL as:

$$x \leftarrow (A[ij] +. \times w[ij]) \div +/ w[ij]$$

©Dennis Bricker, U. of Iowa, 1997