

September 30, 2016


NAME _____

Quiz 5. When a valve is opened, fluid flows in the expansion duct shown below according to the approximation

$$\underline{V} = u\hat{i} = U\left(1 - \frac{x}{2L}\right)\left(\frac{Ut}{L}\right)\hat{i}$$

for $t \ll L/U$. If $L = 1$ m and $U = 1$ m/s, then at $(x, t) = (L, L/2U)$,

- 1) Find the unsteady (local) acceleration of a_x
- 2) Find the convective acceleration of a_x
- 3) Find the total acceleration a_x



Acceleration:

$$a_x = \frac{\partial u}{\partial t} + u \frac{\partial u}{\partial x} + v \frac{\partial u}{\partial y} + w \frac{\partial u}{\partial z}$$

$$a_y = \frac{\partial v}{\partial t} + u \frac{\partial v}{\partial x} + v \frac{\partial v}{\partial y} + w \frac{\partial v}{\partial z}$$

$$a_z = \frac{\partial w}{\partial t} + u \frac{\partial w}{\partial x} + v \frac{\partial w}{\partial y} + w \frac{\partial w}{\partial z}$$

Note: Attendance (+2 points), format (+1 point)

Solution:

- 1) local acceleration

$$(a_x)_{local} = \frac{\partial u}{\partial t} = \frac{\partial}{\partial t} \left[U \left(1 - \frac{x}{2L} \right) \left(\frac{Ut}{L} \right) \right] = \frac{U^2}{L} \left(1 - \frac{x}{2L} \right) \quad (+2 \text{ points})$$

at $(x, t) = (L, L/2U)$,

$$(a_x)_{local} = \frac{U^2}{L} \left(1 - \frac{L}{2L} \right) = \frac{U^2}{2L} = \frac{(1 \text{ m/s})^2}{2 \times 1 \text{ m}} = 0.5 \frac{\text{m}}{\text{s}^2} \quad (+1 \text{ points})$$

- 2) convective acceleration

$$(a_x)_{conv} = u \frac{\partial u}{\partial x} = U \left(1 - \frac{x}{2L} \right) \left(\frac{Ut}{L} \right) \frac{\partial}{\partial x} \left[U \left(1 - \frac{x}{2L} \right) \left(\frac{Ut}{L} \right) \right] = -\frac{U^2}{2L} \left(1 - \frac{x}{2L} \right) \left(\frac{Ut}{L} \right)^2 \quad (+2 \text{ points})$$

at $(x, t) = (L, L/2U)$,

$$(a_x)_{conv} = -\frac{U^2}{2L} \left(1 - \frac{L}{2L} \right) \left(\frac{U}{L} \cdot \frac{L}{2U} \right)^2 = -\frac{U^2}{16L} = -\frac{(1 \text{ m/s})^2}{16 \times 1 \text{ m}} = -0.0625 \frac{\text{m}}{\text{s}^2} \quad (+1 \text{ points})$$

September 30, 2016

3) total acceleration

$$a_x = (a_x)_{local} + (a_x)_{conv} = \frac{\partial u}{\partial t} + u \frac{\partial u}{\partial x} = \frac{U^2}{2L} \left(1 - \frac{x}{2L}\right) \left(1 - \frac{1}{2} \left(\frac{Ut}{L}\right)^2\right) \quad (+0.5 \text{ points})$$

at $(x, t) = (L, L/2U)$,

$$a_x = \frac{U^2}{L} \left(1 - \frac{L}{2L}\right) \left(1 - \frac{1}{2} \left(\frac{U \cdot L}{L \cdot 2U}\right)^2\right) = \frac{7U^2}{16L} = \frac{7 \times (1 \text{ m/s})^2}{16 \times 1 \text{ m}} = 0.4375 \frac{\text{m}}{\text{s}^2} \quad (+0.5 \text{ points})$$